arXiv:1210.6011v1 [math.CV] 22 Oct 2012

THE DYNAMICS OF HOLOMORPHIC CORRESPONDENCES OF $\mathbb{P}^1$: INVARIANT MEASURES AND THE NORMALITY SET

GAUTAM BHARALI AND SHRIHARI SRIDHARAN

Abstract. We investigate whether analogues of Brolin's theorem hold true when iterating a holomorphic correspondence on $\mathbb{P}^1$. Our results have the following character: if F is a holomorphic correspondence on $\mathbb{P}^1$, then (under certain conditions) F admits an invariant measure μ_F such that, for any point z drawn from a "large" open subset of $\mathbb{P}^1$, μ_F is the weak*-limit of suitably normalised sums of point masses carried by the pre-images of z under successive iterates of F. Let ${}^{\dagger}F$ denote the transpose of F. Under the condition $d_{top}(F) > d_{top}({}^{\dagger}F)$, where d_{top} denotes the topological degree, the above is a small refinement of a set of recent results by Dinh and Sibony. Our substantive result, under this condition, is that the support of μ_F is disjoint from the normality set of F. There are many interesting correspondences on $\mathbb{P}^1$ for which $d_{top}(F) \leq d_{top}({}^{\dagger}F)$. Examples are the correspondences introduced by Bullett and collaborators. When $d_{top}(F) \leq d_{top}({}^{\dagger}F)$, equidistribution cannot be expected in general, but we show that something close to classical equidistribution occurs if F has a repeller.

1. Introduction

The dynamics studied in this paper owes its origin to a work of Bullett [2] and to a series of articles motivated by [2] — most notably [5, 6, 4]. The object of study in [2] is the dynamical system that arises on iterating a certain relation on $\mathbb{C}$. This relation is the zero set of a polynomial $g \in \mathbb{C}[z_1, z_2]$ of a certain form such that:

- $g(\cdot, z_2)$ and $g(z_1, \cdot)$ are generically quadratic; and
- no irreducible component of $\{g = 0\}$ is of the form $\{a\} \times \mathbb{C}$ or $\mathbb{C} \times \{a\}$, where $a \in \mathbb{C}$.

The form of g above is such that, if V_g denotes the biprojective completion of $\{g = 0\}$ in $\mathbb{P}^1 \times \mathbb{P}^1$ and π_j denotes the projection onto the jth factor, then the set-valued maps

$$\begin{aligned} \mathbb{P}^1 \ni z_1 &\longmapsto \pi_2\left(\pi_1^{-1}\{z_1\} \cap V_g\right), \quad \text{and} \\ \mathbb{P}^1 \ni z_2 &\longmapsto \pi_1\left(\pi_2^{-1}\{z_2\} \cap V_g\right), \end{aligned} \tag{1.1}$$

are both 2-valued (counting intersections according to multiplicity). In [6], this set-up is extended to polynomials $g \in \mathbb{C}[z_1, z_2]$ of arbitrary degree that induce holomorphic correspondences $V_g \subset \mathbb{P}^1 \times \mathbb{P}^1$ such that the first map given by (1.1) is m-valued and the second map is n-valued, $m, n \geq 2$. It would be interesting to know whether such a correspondence admits a dynamical construction of an invariant measure in analogy to Brolin's Theorem [1, Theorem 16.1]. This problem is the main theme of this paper.

2000 *Mathematics Subject Classification.* Primary 37F05, 37F10; Secondary 32U40.

Key words and phrases. Holomorphic correspondence, invariant measure, normality set.

The first author is supported in part by a UGC Centre of Advanced Study grant.

The reader will be aware of recent results by Dinh and Sibony [9] that, it would seem, should immediately solve the above problem. *However, key assumptions in the theorems of* [9] *fail to hold for many interesting correspondences on* $\mathbb{P}^1$. We shall discuss what this assertion means in the remainder of this section. We will need several definitions to be able to state our results rigorously, which we postpone to Section 3.

Concerning the dynamics of multivalued maps between compact Kähler manifolds: results of perhaps the broadest scope are established in [9]. In the context of correspondences, we borrow from this work our terminology and the following definition.

Definition 1.1. Let X_1 and X_2 be two compact complex manifolds of dimension k. We say that $\mathit{\Gamma}$ is a *holomorphic k-chain* in $X_1 \times X_2$ if $\mathit{\Gamma}$ is a formal linear combination of the form

$$\mathit{\Gamma} = \sum_{j=1}^{N} m_j \Gamma_j, \tag{1.2}$$

where the m_j's are positive integers and $\Gamma_1, \dots, \Gamma_N$ are distinct irreducible complex subvarieties of $X_1 \times X_2$ of pure dimension k. Let π_i denote the projection onto X_i, $i = 1, 2$. We say that the holomorphic k-chain $\mathit{\Gamma}$ *gives a meromorphic correspondence of* X_1 *onto* X_2 if, for each Γ_j in (1.2), $\pi_1|_{\Gamma_j}$ and $\pi_2|_{\Gamma_j}$ are surjective. The set-valued map, denoted as $F_{\mathit{\Gamma}}$, given by

$$X_1 \ni x \longmapsto \bigcup_{j=1}^{N} \pi_2\left(\pi_1^{-1}\{x\} \cap \Gamma_j\right)$$

is called the *meromorphic correspondence determined by* $\mathit{\Gamma}$. We call $F_{\mathit{\Gamma}}$ a *holomorphic correspondence* if $F_{\mathit{\Gamma}}(x)$ is a finite set for every $x \in X_1$.

Remark 1.2. It is helpful to encode meromorphic correspondences as holomorphic chains. We will encounter circumstances where, in the notation of (1.2), $m_j \geq 2$. For instance: even if we start with a holomorphic correspondence on $\mathbb{P}^1$ determined by the biprojective completion, say $V_g \subset \mathbb{P}^1 \times \mathbb{P}^1$, of an *irreducible* polynomial $g \in \mathbb{C}[z_1, z_2]$, composing V_g with itself (see Section 2) can result in a variety that is not irreducible and some of whose irreducible components occur with multiplicity greater than 1.

Suppose (X, ω) is a compact Kähler manifold of dimension k (ω denoting the normalised Kähler form) and F is a meromorphic correspondence of X onto itself. One of the results in [9] says, roughly, that if $d_{k-1}(F) < d_k(F)$, where $d_{k-1}(F)$ and $d_k(F)$ are the dynamical degrees of F of order $(k-1)$ and k respectively, then there exists a probability measure μ_F satisfying $F^*\mu_F = d_k(F)\mu_F$, such that

$$\frac{1}{d_k(F)^n}(F^n)^*(\omega^k) \xrightarrow{\text{weak}^*} \mu_F \text{ as measures, as } n \to \infty. \tag{1.3}$$

When $\dim_{\mathbb{C}}(X) = 1$, the assumption $d_{k-1}(F) < d_k(F)$ translates into the assumption that the (generic) number of pre-images under F is *strictly larger* than the number of images under F, both counted according to multiplicity. One of the important applications of the study of the iterates of meromorphic correspondences is to study the dynamics of dominant meromorphic self-maps of X with $\dim_{\mathbb{C}}(X) \geq 2$. In this context, the assumption $d_{k-1}(F) < d_k(F)$ is a very natural one. However, none of the techniques in the current literature are of help in studying correspondences F for which $d_{k-1}(F) \geq d_k(F)$, even when $(X,\omega) = (\mathbb{P}^k, \omega_{FS})$ and meaningful asymptotics are expected (in this paper ω_{FS} will denote the Kähler form associated to the Fubini–Study metric).

Why should one be interested in the dynamics of a meromorphic correspondence $F : X \to X$, X a compact Kähler manifold, for which $d_{k-1}(F) \geq d_k(F)$? The work of Bullett and collaborators suggest several reasons in the case $(X,\omega) = (\mathbb{P}^1, \omega_{FS})$. Thus, *we shall focus on correspondences on* $\mathbb{P}^1$ (although parts of our results hold true for Riemann surfaces). A *mating* of two monic polynomials on $\mathbb{C}$ is a construction by Douady [11] that, given two monic polynomials $f, g \in \mathbb{C}[z]$ of the same degree, produces a continuous branched covering $(f \amalg g)$ of the topological sphere to itself. For certain natural choices of pairs (f, g), one can determine in principle — see [13, Theorem 2.1] — when $(f \amalg g)$ is semiconjugate to a rational map on $\mathbb{P}^1$. In a series of papers by Bullett and collaborators [5, 3, 4], these ideas have been extended to matings between polynomials and certain discrete subgroups of the Möbius group or certain Hecke groups, whose dynamics turn out to be conjugate to the dynamics of holomorphic correspondences on $\mathbb{P}^1$. To be a bit more precise: in these constructions, $\mathbb{P}^1$ is partitioned into two regions restricted to which the action of some holomorphic correspondence on $\mathbb{P}^1$ is conjugate to either the (iterative) action of a monic polynomial on its filled Julia set or to the standard action of one of the groups just mentioned on some component of its regular set. It would be interesting to build an ergodic theory for such matings. Now, *in all the known constructions where a holomorphic correspondence* $F : \mathbb{P}^1 \to \mathbb{P}^1$ *models the dynamics of a mating of some polynomial with some group,* $d_0(F) = d_1(F)$. Therefore, in this context, producing an invariant measure (and doing so via iterated inverse images) would require that the techniques in [9] be supplemented by other ideas.

Before turning to a rigorous statement of our results, we give an informal description of our work. First: when F is a holomorphic correspondence on $\mathbb{P}^1$, and $d_0(F) < d_1(F)$, there exists a polar set $\mathcal{E} \subsetneq \mathbb{P}^1$ such that

$$\frac{1}{d_1(F)^n}(F^n)^*(\delta_x) \xrightarrow{\text{weak}^*} \mu_F \text{ as measures, as } n \to \infty, \ \forall x \in \mathbb{P}^1 \setminus \mathcal{E}, \tag{1.4}$$

where μ_F is as in (1.3) with $(X,\omega) = (\mathbb{P}^1, \omega_{FS})$. This follows almost directly from (1.3) and another result in [9]. But furthermore, we shall see — and *this* is the substantive half of our result — that the support of μ_F is disjoint from the normality set of F, where "normality set" is a notion of the Fatou set introduced in [6] in the context of correspondences.

When F (a holomorphic correspondence on $\mathbb{P}^1$) satisfies $d_0(F) \geq d_1(F)$, there is no reason to expect (1.4). Indeed, consider the examples: $F_1(z) := 1/z$, in which case $d_0(F_1) = d_1(F_1) = 1$, or F_2 being the holomorphic correspondence whose graph is the zero set in $\mathbb{P}^1 \times \mathbb{P}^1$ of the polynomial $g(z_1, z_2) = z_2^2 - (z_1^2 + 1)$, in which case $d_0(F_2) = d_1(F_2) = 2$. When $d_0(F) \geq d_1(F)$, we draw upon certain ideas of McGehee [15]. We find that if F admits a repeller — in the sense of McGehee — $\mathcal{R} \subset \mathbb{P}^1$ having certain properties, then there exists a neighbourhood $U(F,\mathcal{R}) \supset \mathcal{R}$ and a probability measure μ_F satisfying $F^*\mu_F = d_1(F)\mu_F$, such that

$$\frac{1}{d_1(F)^n}(F^n)^*(\delta_x) \xrightarrow{\text{weak}^*} \mu_F \text{ as measures, as } n \to \infty, \ \forall x \in U(F,\mathcal{R}). \tag{1.5}$$

A rigorous statement of this is given by Theorem 3.4. Note that (1.5) differs from (1.4) in that it does not state that $\mathbb{P}^1 \setminus U(F, \mathcal{R})$ is polar (or even nowhere dense), but this is the best one can expect (see Remark 3.5 below).

2. Fundamental definitions

In this section, we isolate certain definitions that are *fundamental* to our results but are somewhat involved. Since our results pertain to correspondences on $\mathbb{P}^1$, on which (for dimensional reasons) meromorphic correspondences are automatically holomorphic, we shall focus *only* on holomorphic correspondences.

2.1. **The composition of two holomorphic correspondences.**

Let X be a complex manifold of dimension k. For any holomorphic k-chain $\mathit{\Gamma}$ on X, we define the *support* of $\mathit{\Gamma}$, assuming the representation (1.2), by

$$|\mathit{\Gamma}| := \cup_{j=1}^{N} \Gamma_j.$$

Consider two holomorphic correspondences, determined by the k-chains

$$\mathit{\Gamma}^1 = \sum_{j=1}^{N_1} m_{1,j}\Gamma_{1,j}, \qquad \mathit{\Gamma}^2 = \sum_{j=1}^{N_2} m_{2,j}\Gamma_{2,j},$$

in $X \times X$. The k-chains $\mathit{\Gamma}_1$, $\mathit{\Gamma}_2$ have the alternative representations

$$\mathit{\Gamma}^1 = {\sum}'_{1\leq j\leq L_1} \Gamma^{\bullet}_{1,j}, \qquad \mathit{\Gamma}^2 = {\sum}'_{1\leq j\leq L_2} \Gamma^{\bullet}_{2,j}, \tag{2.1}$$

where the primed sums indicate that, in the above representation, the irreducible subvarieties $\Gamma^{\bullet}_{s,j}$, $j = 1, \dots, L_s$, $s = 1, 2$, are *not necessarily distinct* and are repeated according to the coefficients $m_{s,j}$. We define the holomorphic k-chain $\Gamma^{\bullet}_{2,l} \circ \Gamma^{\bullet}_{1,j}$ by the following two requirements:

$$|\Gamma^{\bullet}_{2,l} \circ \Gamma^{\bullet}_{1,j}| := \{(x, z) \in X \times X : \exists y \in X \text{ s.t. } (x, y) \in \Gamma^{\bullet}_{1,j},\ (y, z) \in \Gamma^{\bullet}_{2,l}\}, \tag{2.2}$$

$$\Gamma^{\bullet}_{2,l} \circ \Gamma^{\bullet}_{1,j} \equiv \sum_{1\leq s\leq N(j,l)} \nu_{s,jl} Y_{s,jl},$$

where the $Y_{s,jl}$'s are the distinct irreducible components of the subvariety on the right-hand side of (2.2), and $\nu_{s,jl} \in \mathbb{Z}_+$ is the generic number y's as (x, z) varies through $Y_{s,jl}$ for which the membership conditions on the right-hand side of (2.2) are satisfied. Finally, we define the k-chain

$$\mathit{\Gamma}^2 \circ \mathit{\Gamma}^1 := \sum_{j=1}^{L_1} \sum_{l=1}^{L_2} \Gamma^{\bullet}_{2,l} \circ \Gamma^{\bullet}_{1,j}. \tag{2.3}$$

If either $\mathit{\Gamma}^1$ or $\mathit{\Gamma}^2$ is a meromorphic, *non-holomorphic* correspondence, then the composition rule is *considerably more technical* than given by (2.2); see [9]. The definition in [9] is required to ensure that no $\Gamma^{\bullet}_{2,l} \circ \Gamma^{\bullet}_{1,j}$ has irreducible components whose projections are not equal to X, and is equivalent to (2.2) when both $\mathit{\Gamma}^1$ and $\mathit{\Gamma}^2$ are holomorphic — see, for example, [10, Section 4]. The last fact requires a certain amount of intersection theory, but the interested reader is directed to Section 4 for an *elementary* proof of this fact when $\mathit{\Gamma}^1$ and $\mathit{\Gamma}^2$ are holomorphic correspondences on $\mathbb{P}^1$. To summarise: if $\mathit{\Gamma}^1$ and

$\mathit{\Gamma}^2$ determine holomorphic correspondences on X, then so does $\mathit{\Gamma}^2 \circ \mathit{\Gamma}^1$. The n-fold iterate of $\mathit{\Gamma}$ will be denoted by $\mathit{\Gamma}^{\circ n}$.

Given a k-chain $\mathit{\Gamma}$ on X (X as above), we may view $|\mathit{\Gamma}|$ as a relation of X to itself. In certain sections of this paper, when we specialise to $X = \mathbb{P}^1$, we will need to make *essential* use of McGehee's results from [15] on the dynamics of closed relations on compact spaces. Let $\mathit{\Gamma}^1$ and $\mathit{\Gamma}^2$ be as above. Then, the composition of the two relations $|\mathit{\Gamma}^1|$ and $|\mathit{\Gamma}^2|$ in the classical sense (denoted here by $|\mathit{\Gamma}^2| \star |\mathit{\Gamma}^1|$) — which is the sense in which the term is used in [15] — is defined as

$$|\mathit{\Gamma}^2| \star |\mathit{\Gamma}^1| \ := \ \{(x,z) \in X \times X : \exists y \in X \text{ such that } (x,y) \in |\mathit{\Gamma}^1|, \ (y,z) \in |\mathit{\Gamma}^2|\}. \tag{2.4}$$

It is easy to see that for holomorphic correspondences

$$|\mathit{\Gamma}^2 \circ \mathit{\Gamma}^1| \ = \ |\mathit{\Gamma}^2| \star |\mathit{\Gamma}^1|. \tag{2.5}$$

In what follows, we shall adopt a notational simplification. Given a k-chain that determines a holomorphic correspondence, and there is no scope for confusion, we shall simply denote $F_{\mathit{\Gamma}}$ by F.

2.2. **The normality set of a holomorphic correspondence on a Riemann surface.** Let F be a holomorphic correspondence on a compact Riemann surface X. The motivation for the concept of the *normality set of* F is quite simple. However, we will need a formalism that will enable good book-keeping. In this subsection, we will use the representation (2.1) for a holomorphic correspondence $\mathit{\Gamma}$. The set of integers $\{m, m+1, \dots, n\}$ will be denoted by $[m\,..\,n]$.

Given $N \in \mathbb{Z}_+$, we say that $(z_0, \dots, z_N; \, \alpha_1, \dots, \alpha_N) \in X^{N+1} \times [1\,..\,L]^N$ (see (2.1) for the meaning of L) is *a path of an iteration of* F *starting at* z_0*, of length* N, or simply an N*-path starting at* z_0, if

$$(z_{j-1}, z_j) \in \Gamma^{\bullet}_{\alpha_j}, \ \ j = 1, \dots, N.$$

Next, given any two irreducible subvarieties $\Gamma^{\bullet}_j$ and $\Gamma^{\bullet}_k$ in the decomposition of $\mathit{\Gamma}$ in the sense of (2.1), we define

$$\begin{aligned}\Gamma^{\bullet}_{(j,k)} \ \equiv \ \Gamma^{\bullet}_k \wedge \Gamma^{\bullet}_j \ &:= \{(x,y,z) \in X^3 : (x,y) \in \Gamma^{\bullet}_j, \ (y,z) \in \Gamma^{\bullet}_k\} \\ &= (\Gamma^{\bullet}_j \times X) \cap (X \times \Gamma^{\bullet}_k).\end{aligned}$$

For any multi-index $\boldsymbol{\alpha} \in [1\,..\,L]^j$, $\boldsymbol{\alpha} = (\alpha_1, \dots, \alpha_j)$, we recursively define

$$\Gamma^{\bullet}_{\boldsymbol{\alpha}} \ := \ (\Gamma^{\bullet}_{(\alpha_1,\dots,\alpha_{j-1})} \times X) \cap (X^{j-1} \times \Gamma^{\bullet}_{\alpha_j}) \ \text{ for } j \geq 2. \tag{2.6}$$

In all discussions on the normality set of F, we shall assume the following for any N-path $(z_0, \dots, z_N; \, \alpha_1, \dots, \alpha_N)$:

($*$) *For each* $j = 1, \dots, N$, $\Gamma^{\bullet}_{(\alpha_1,\dots,\alpha_j)} \cap B_j$ *is an irreducible subvariety of* B_j *for every sufficiently small ball* $B_j \ni (z_0, \dots, z_j)$.

An N-path will be called an *admissible* N*-path* if it satisfies ($*$). Now fix a $z_0 \in X$ and $N \in \mathbb{Z}_+$, and set

$$\mathscr{P}_N(F, z_0) \ := \ \text{the set of all paths of iterations of } F\text{, of length } N\text{, starting at } z_0.$$

We will denote an element of $\mathscr{P}_N(F, z_0)$ either by $\boldsymbol{\mathscr{Z}}$ or by $(Z; \, \boldsymbol{\alpha}) \in X^{N+1} \times [1\,..\,L]^N$, depending on the need. Observe that if $\boldsymbol{\mathscr{Z}}$ is an admissible N-path, $N \geq 2$, then there is a *unique* irreducible component of $\Gamma^\bullet_{(\alpha_1,\dots,\alpha_j)}$ to which $(z_0,\dots,z_j)$ belongs, $j=2,\dots,N$. Hence, if $\boldsymbol{\mathcal{Z}}$ is an admissible N-path, let us write

$$\Gamma^\bullet_{\boldsymbol{\mathcal{Z}},j} := \begin{cases} \Gamma^\bullet_{\alpha_1}, & \text{if } j=1,\\ \text{the unique irreducible component of } \Gamma^\bullet_{(\alpha_1,\dots,\alpha_j)} \text{ containing } (z_0,\dots,z_j), & \text{if } j\geq 2. \end{cases}$$

Finally, let $\nu_{(\boldsymbol{\mathcal{Z}},j)} : Y_{\boldsymbol{\mathcal{Z}},j} \longrightarrow \Gamma^\bullet_{\boldsymbol{\mathcal{Z}},j}$, where $Y_{\boldsymbol{\mathcal{Z}},j}$ is a compact Riemann surface, denote the desingularization of $\Gamma^\bullet_{\boldsymbol{\mathcal{Z}},j}$. We now have the essential notations needed to define the normality set — in the sense of Bullett–Penrose [6] — of the holomorphic correspondence F. *The one departure that we make from the Bullett–Penrose definition* is that, in defining a holomorphic branch of an iteration along $\boldsymbol{\mathcal{Z}}$, we assume that $\boldsymbol{\mathcal{Z}}$ is admissible. The only purpose of this restriction is that, in proving Theorem 3.1-(*ii*), we will require precise book-keeping of all the branches involved, but *we do not want the simple motivation for the normality set to be obscured by too much book-keeping paraphernalia.*

The motivation for the normality set of F is quite simple: it is the analogue of the Fatou set. However, some additional formalism is required to achieve this analogy in view of the following essential considerations:

- *a*) The analogue for the family of all iterates of a map must be the collection of local maps defined by the germ of $\Gamma^{\circ N}$ at (z_0, z_N), *whenever this makes sense* (around a chosen z_0) for every $\boldsymbol{\mathcal{Z}} \in \mathscr{P}_N(F, z_0)$ and for every $N \in \mathbb{Z}_+$.
- *b*) Since the germs in (*a*) might have singularities, we must pass to Puiseux parametrizations to identify the family of functions (defined around z_0) that we require to be normal.

Let proj_k and π^j_k denote the following projections:

$$\mathsf{proj}_k : X^{k+1} \longrightarrow X^k,\ \ \mathsf{proj}_k : (x_0,\dots,x_k) \longmapsto (x_0,\dots,x_{k-1}),$$
$$\pi^j_k : X^{k+1} \longrightarrow X,\ \ \pi^j_k : (x_0,\dots,x_k) \longmapsto x_j,\ \ 0\leq j\leq k,$$

where $k \in \mathbb{Z}_+$. We now have all the notations for the key definition needed on the way to defining the normality set of F. A schematic drawing is presented on the following page to give an impression of the various objects occurring in the following definition.

Definition 2.1. Let X be a compact Riemann surface and let F be a holomorphic correspondence on X. Let $N \in \mathbb{Z}_+$, $z_0 \in X$, and let $\boldsymbol{\mathcal{Z}} \in \mathscr{P}_N(F, z_0)$. Write $\boldsymbol{\mathcal{Z}} = (Z; \boldsymbol{\alpha}) = (z_0,\dots,z_N; \boldsymbol{\alpha})$. We call the list $(\nu_{(\boldsymbol{\mathcal{Z}},1)} \circ \psi_{(\boldsymbol{\mathcal{Z}},1)},\dots,\nu_{(\boldsymbol{\mathcal{Z}},N)} \circ \psi_{(\boldsymbol{\mathcal{Z}},N)}; U)$ a *holomorphic branch of an iteration of F along $\boldsymbol{\mathcal{Z}}$* if $\boldsymbol{\mathcal{Z}}$ is an admissible N-path, U is a connected neighbourhood of z_0 and $\psi_{(\boldsymbol{\mathcal{Z}},j)} : D \longrightarrow Y_{(\boldsymbol{\mathcal{Z}},j)}$ are holomorphic mappings defined on a planar domain D containing 0 such that, for each $j = 1,\dots,N$:

1) $\nu_{(\boldsymbol{\mathcal{Z}},j)} \circ \psi_{(\boldsymbol{\mathcal{Z}},j)}(0) = (z_0,\dots,z_j)$.
2) The set $U(Z,\boldsymbol{\alpha},j) :=$ the irreducible component of $\Omega(Z,\boldsymbol{\alpha},j) \cap \Gamma^\bullet_{\boldsymbol{\mathcal{Z}},j}$ containing $(z_0,\dots,z_j)$ $\big($where we define

$$\Omega(Z,\boldsymbol{\alpha},j) := \begin{cases} \mathsf{proj}_1^{-1}(U), & \text{if } j=1,\\ \mathsf{proj}_j^{-1}\left[U \times \prod_{k=1}^{j-1} \pi^k_k \circ \nu_{(\boldsymbol{\mathcal{Z}},k)} \circ \psi_{(\boldsymbol{\mathcal{Z}},k)}(D)\right], & \text{if } j\geq 2\,\big) \end{cases}$$

is such that $\pi^j_j\big|_{U(Z,\boldsymbol{\alpha},j)}$ is a (perhaps ramified) covering map.


$\Omega(Z,\boldsymbol{\alpha},2)$
$U(Z,\boldsymbol{\alpha},2)$
$\Gamma^{\bullet}_{\boldsymbol{\mathcal{Z}},2}$
$\nu_{(\boldsymbol{\mathcal{Z}},\alpha,2)}$
(z_0,z_1,z_2)
$\nu^{-1}_{(\boldsymbol{\mathcal{Z}},\alpha,2)}\{p\}$
p
$\nu^{-1}_{(\boldsymbol{\mathcal{Z}},\alpha,2)}(U(Z,\boldsymbol{\alpha},2))$
z_0
$\Gamma^{\bullet}_1$
$\Gamma^{\bullet}_2$
z_1
$\Omega(Z,\boldsymbol{\alpha},1)$
$Y_{(\boldsymbol{\mathcal{Z}},2)}$
$X\times X\times X$

FIGURE 1. Objects associated to Definition 2.1; $\alpha_1 = 1$; $\Gamma^{\bullet}_{\boldsymbol{\mathcal{Z}},2}$ partially obscured by $\Omega(Z,\boldsymbol{\alpha},2)$

3) $\psi_{(\boldsymbol{\mathcal{Z}},j)}$ is a finite-sheeted (perhaps ramified) covering of $\nu^{-1}_{(\boldsymbol{\mathcal{Z}},j)}(U(Z,\boldsymbol{\alpha},j))$.
4) $\mathsf{proj}_j \circ \nu_{(\boldsymbol{\mathcal{Z}},j)} \circ \psi_{(\boldsymbol{\mathcal{Z}},j)} = \nu_{(\boldsymbol{\mathcal{Z}},j-1)} \circ \psi_{(\boldsymbol{\mathcal{Z}},j-1)}$, provided $j \neq 1$.

Remark 2.2. As $\boldsymbol{\mathcal{Z}}$ is admissible, $U(Z,\boldsymbol{\alpha},j)$ in condition (2) is well-defined and $\nu^{-1}_{(\boldsymbol{\mathcal{Z}},j)}(U(Z,\boldsymbol{\alpha},j))$ is a smooth patch that parametrises it; see [12, Chapter 7].

Note that if F is a non-constant rational map on $\mathbb{P}^1$ then

$$\nu_{(\boldsymbol{\mathcal{Z}},j)} \circ \psi_{(\boldsymbol{\mathcal{Z}},j)} \;:=\; \left(\mathsf{id}_D + z_0,\, F(\cdot + z_0)|_D\,, \dots,\, F^j(\cdot + z_0)\big|_D\right), \quad j = 1,2,3,\dots,$$

where D is a small disc around 0, satisfy all the conditions in the above definition. This is the situation that Definition 2.1 generalises. This definition might look very different from that given in [6], but that is largely because we need *more explicit* labelling of the various holomorphic branches than what the notations in [6] provide. It is in this effort of labelling that the assumption $(*)$ helps. Now, having defined holomorphic branches, we can give the following definition.

Definition 2.3 (paraphrasing Bullet–Penrose, [6])**.** Let X be a compact Riemann surface and let F be a holomorphic correspondence on X. A point $z_0 \in X$ is said to belong to the *normality set of* F, denoted by $\mathscr{N}(F)$, if there exists a connected neighbourhood U of z_0 and a *single* planar domain D containing 0, which depends on z_0, such that

1) For each $N \in \mathbb{Z}_+$ and each $\boldsymbol{\mathscr{Z}} \in \mathscr{P}_N(F, z_0)$, there exists a holomorphic branch $(\nu_{(\boldsymbol{\mathscr{Z}},1)} \circ \psi_{(\boldsymbol{\mathscr{Z}},1)}, \dots, \nu_{(\boldsymbol{\mathscr{Z}},N)} \circ \psi_{(\boldsymbol{\mathscr{Z}},N)}; U)$ of an iteration of F along $\boldsymbol{\mathscr{Z}}$ with $\mathsf{Dom}(\psi_{(\boldsymbol{\mathscr{Z}},j)}) = D$ for every $(\boldsymbol{\mathscr{Z}}, j)$.
2) The family
$$\begin{aligned}\mathscr{F}(z_0) \ :=\ \big\{\pi^N_N \circ \nu_{(\boldsymbol{\mathscr{Z}},N)} \circ \psi_{(\boldsymbol{\mathscr{Z}},N)} : N \in \mathbb{Z}_+,\ &\boldsymbol{\mathscr{Z}} \in \mathscr{P}_N(F, z_0), \text{ and} \\ &(\nu_{(\boldsymbol{\mathscr{Z}},1)} \circ \psi_{(\boldsymbol{\mathscr{Z}},1)}, \dots, \nu_{(\boldsymbol{\mathscr{Z}},N)} \circ \psi_{(\boldsymbol{\mathscr{Z}},N)}; U) \text{ is a holomorphic} \\ &\ \text{branch of an iteration of } F \text{ along } \boldsymbol{\mathscr{Z}}\big\}\end{aligned}$$
is a normal family on D.

Remark 2.4. The set $\mathscr{N}(F)$ is open, although it is not necessarily non-empty. If $z_0 \in \mathscr{N}(F)$ and U is the neighbourhood of z_0 as given by Definition 2.3, then it is routine to show that $U \subset \mathscr{N}(F)$.

3. More definitions and statement of results

We need to present some formalisms before we can state our first result.

Given a holomorphic correspondence on X, $\dim_{\mathbb{C}}(X) = k$, determined by a holomorphic k-chain $\varGamma$, its *adjoint correspondence* is the meromorphic correspondence determined by the k-chain (assuming the representation (1.2) for $\varGamma$)

$$\begin{aligned} {}^{\dagger}\varGamma \ &:=\ \sum_{j=1}^{N} m_j\, {}^{\dagger}\Gamma_j, \\ \text{where } {}^{\dagger}\Gamma_j \ &:=\ \{(y, x) \in X \times X : (x, y) \in \Gamma_j\}. \end{aligned}$$

In the abbreviated notation introduced in Section 2, we shall henceforth write:

$$F^n \ :=\ F_{\varGamma^{\circ n}}, \qquad {}^{\dagger}F \ :=\ F_{{}^{\dagger}\varGamma}.$$

Given a holomorphic k-chain $\varGamma$ on $X \times X$, $\varGamma$ detemines a current of bidimension (k, k) via the currents of integration given by its constituent subvarieties Γ_j. We denote this current by $[\varGamma]$. If F is the meromorphic correspondence determined by $\varGamma$, we *formally* define the action of F on currents S on X of bidegree (p, p), $0 \leq p \leq k$, by the prescription:

$$F^*(S) \ :=\ (\pi_1)_* \left(\pi_2^*(S) \wedge [\varGamma]\right), \tag{3.1}$$

where, as usual, π_j denotes the projection of $X \times X$ onto the jth factor. This prescription would make sense for those currents for which the pullback by π_2 makes sense and the intersection of this new current with $[\varGamma]$ also makes sense. That this is the case is easy to see when S is a smooth (p, p) form (hence a current of bidegree (p, p) on X). The reader is referred to [9, Section 2.4] for details.

It turns out that if μ is a finite, positive Borel measure, then the intersection with $[\varGamma]$ in (3.1) makes sense. Here, μ is viewed as a current of bidegree (k, k). Let us work out

$F^*\mu$ for a specific example that is central to this paper. Let $x \in X$ and let δ_x be the Dirac mass at x. The prescription (3.1) is interpreted as

$$\begin{aligned}\langle F^*(\delta_x), \varphi\rangle \;=\; \langle \pi_2^*(\delta_x) \wedge [\mathit{\Gamma}], \pi_1^*\varphi\rangle \;&:=\; \sum_{j=1}^{N} m_j \langle (\pi_2|_{\Gamma_j})^*(\delta_x), \pi_1^*\varphi\rangle \\ &=\; \sum_{j=1}^{N} m_j \langle \delta_x, \big(\,\pi_2|_{\Gamma_j}\big)_*(\pi_1^*\varphi)\rangle, \qquad (3.2)\end{aligned}$$

where the last equality is the just the way we define the pullback of a current under a mapping (of an analytic space) that is submersive on a Zariski open subset of its domain. How does one compute the expression (3.2) ? If $\Omega \subset X$ is a Zariski-open subset of X such that $(\pi_2^{-1}(\Omega) \cap \Gamma_j, \Omega, \pi_2)$ is a covering space for each $j = 1, \dots, N$, then, for $x \in \Omega$, $(\pi_2|_{\Gamma_j})_*(f)(x)$ is just the sum of the values of f on the fibre $\pi_2^{-1}\{x\} \cap \Gamma_j$ for any $f \in \mathcal{C}^\infty(X \times X)$. Thus, when $x \in \Omega$, (3.2) equals the quantity

$$\sum_{j=1}^{N} m_j \sum_{\zeta:(\zeta,x)\in\Gamma_j} \varphi(\zeta) \;=:\; \mathbf{\Lambda}[\varphi](x) \qquad x \in \Omega. \qquad (3.3)$$

Now, *if F is holomorphic*, then for any fixed *continuous* function φ, $\mathbf{\Lambda}[\varphi]$ extends continuously to each $x \in X \setminus \Omega$. We shall denote this continuous extension of the left-hand side of (3.3) also as $\mathbf{\Lambda}[\varphi]$. In other words, $F^*(\delta_x)$ can be defined as a measure supported on the set ${}^{\dagger}F(x)$, and

$$\langle F^*(\delta_x), \varphi\rangle \;=\; \mathbf{\Lambda}[\varphi](x) \;\; \forall x \in X, \;\; \forall \varphi \in \mathcal{C}(X). \qquad (3.4)$$

The arguments preceding (3.3) continue to be valid if, in (3.2), δ_x is replaced by μ, a finite, positive Borel measure on X.

The push-forward of a current S by F is defined by the equation $F_*(S) := {}^{\dagger}F^*(S)$ whenever the latter makes sense.

If μ is a Borel measure, the notion that μ is invariant under a correspondence requires appropriate modifications. As before, let X be a compact complex manifold and F a meromorphic correspondence as in (1.2). A Borel measure μ on X is said to be *invariant under X* if

$$\sum_{j=1}^{N} m_j\, \mu(\pi_1(\pi_2^{-1}(E) \cap \Gamma_j)) \;=\; \mu(E) \quad \text{for each Borel set } E \subset X.$$

Note that the sets occurring in the left-hand side of the above equation are Borel sets.

Finally, we define two numbers that are essential to the statement of our theorems. With F as above, let $d_1(F)$ denote the generic number of preimages under F of a point in $\mathbb{P}^1$, counted according to multiplicity. What we mean by "counted according to multiplicity" is that if Ω is the Zariski-open set defined prior to the equation (3.3), and the 1-chain $\mathit{\Gamma}$ determining F has the representation (1.2), then

$$d_1(F) \;=\; \sum_{j=1}^{N} m_j \mathsf{Card}\{x : (x,y) \in \Gamma_j\}, \quad y \in \Omega,$$

irrespective of the choice of $y \in \Omega$. In other words, $d_1(F)$ is the *topological degree of* F, denoted as $d_{top}(F)$. Define $d_0(F) := d_{top}({}^{\dagger}F)$. With these definitions, we are ready to state our first theorem.

Theorem 3.1. *Let* F *be a holomorphic correspondence on* $\mathbb{P}^1$ *and assume that* $d_0(F) < d_1(F)$. *Then:*

i) *There exist a Borel probability measure* μ_F *on* $\mathbb{P}^1$ *that satisfies* $F^*(\mu_F) = d_1(F)\mu_F$ *and a polar set* $\mathcal{E} \subsetneq \mathbb{P}^1$ *such that for each* $z \in \mathbb{P}^1 \setminus \mathcal{E}$

$$\frac{1}{d_1(F)^n}(F^n)^*(\delta_z) \xrightarrow{\textit{weak}^*} \mu_F \;\; \textit{as measures, as } n \to \infty.$$

In particular, μ_F *is invariant under* F.

ii) $\operatorname{supp}(\mu_F)$ *is disjoint from the normality set of* F.

The proof of the above theorem may be summarised as follows.

- Since $d_0(F) < d_1(F)$, a direct application of two results from [9] is possible, from which it follows that $d_1(F)^{-n}(F^n)^*(\delta_z) \longrightarrow \mu_F$ in the sense of distributions.
- One can show that, owing to dimensional considerations, convergence in the sense of distributions in the present set-up coincides with weak* convergence in $\mathcal{C}(\mathbb{P}^1;\mathbb{R})^*$.
- To establish Part (*ii*) we appeal a second time to one of the results used to establish Part (*i*) — i.e. to (1.3) with $\omega = \omega_{FS}$. Given this, it suffices to show that the masses of any compact $K \Subset \mathcal{N}(F)$ with respect to the measures induced by $(F^n)^*(\omega_{FS})$ are bounded from above.

The situation is *very different* when $d_0(F) \geq d_1(F)$. To repeat: we should not expect asymptotic equidistribution of preimages in general, even when $d_0(F), d_1(F) \geq 2$, as the holomorphic correspondence F whose graph is the zero set in $\mathbb{P}^1 \times \mathbb{P}^1$ of the polynomial $g(z_1, z_2) = z_2^2 - (z_1^2 + 1)$ (where $d_0(F) = d_1(F) = 2$) illustrates. We require some dynamically meaningful condition for things to work. It is this need that motivates the next few definitions. Let X be a compact Hausdorff space and let $f \subset X \times X$ be a relation of X to itself such that $\pi_1(f) = X$. For any set $S \subset X$, we write

$$f(S) \; := \; \pi_2\left(\pi_1^{-1}(S) \cap f\right).$$

We define the nth iterated relation by

$$\begin{aligned} f^{(n)} \; &:= \; f \star f^{(n-1)} \;\; \text{for } n \geq 2, \\ f^{(1)} \; &:= \; f, \end{aligned} \tag{3.5}$$

where the composition operation $\star$ is as understood from (2.4) above. It is useful to have a notion of omega limit sets in the context of iterating a relation analogous to the case of maps. This definition is provided by McGehee in [15, Section 5]. Following McGehee, for a subset $S \subset X$, let us write

$$\mathfrak{K}(S; f) \; := \{K \subset_{\text{closed}} X : K \text{ satisfies } f(K) \subset K \text{ and } f^{(n)}(S) \subset K \text{ for some } n \geq 0\}$$

(with the understanding that $f^{(0)}$ is the diagonal in $X \times X$). The *omega limit set of* S *under* f, denoted by $\omega(S; f)$, is the set

$$\omega(S; f) \; := \; \bigcap \mathfrak{K}(S; f).$$

We say that a set $\mathcal{A} \subset X$ is an *attractor* for f if $\mathcal{A} \neq X$ and there exists a set U such that $\mathcal{A} \subset U^\circ$ and such that $\omega(U; f) = \mathcal{A}$.

These concepts motivate the following two definitions in the case of holomorphic correspondences.

Definition 3.2. Let F be a holomorphic correspondence on a Riemann surface X given by the holomorphic 1-chain Γ. A set $\mathcal{A} \subset X$ is called an *attractor* for F if it is an attractor for the relation $|\Gamma|$ in the sense of [15]. A set $\mathcal{R}$ is called a *repeller* for F if it is an attractor for the relation $|{}^\dagger\Gamma|$.

We must note here McGehee calls the relation on X induced by $|{}^\dagger\Gamma|$ the *transpose* of $|\Gamma|$, and our $|{}^\dagger\Gamma|$ is $|\Gamma|^*$ in the notation of [15].

Definition 3.3. Let F be as above and let $\mathcal{R}$ be a repeller for F. We say that $\mathcal{R}$ is a *strong repeller* for F if there exists a point $a_0 \in \mathcal{R}$ and an open neighbourhood U of $\mathcal{R}$ such that for each $w \in U$, there exists a sequence $\{a_n(w)\}_{n\in\mathbb{Z}_+}$ such that

- $a_n(w) \in ({}^\dagger F)^n(w)\ \forall n \in \mathbb{Z}_+$; and
- $a_n(w) \longrightarrow a_0$ as $n \to \infty$.

Given a holomorphic correspondence F on $\mathbb{P}^1$ determined by the 1-chain Γ, we write

$$F(S) \ := \ \cup_{j=1}^{N} \pi_2\left(\pi_1^{-1}(S) \cap \Gamma_j\right) \tag{3.6}$$

for any $S \subset \mathbb{P}^1$ (assuming the representation (1.2) for Γ). We call $w \in \mathbb{P}^1$ a *critical value* if there exists an irreducible component Γ_j such that at least one of the irreducible germs of Γ_j at some point in $\pi_2^{-1}\{w\} \cap \Gamma_j$ is either non-smooth or does *not* project injectively under π_2.

We are now in a position to state our next result.

Theorem 3.4. *Let F be a holomorphic correspondence on $\mathbb{P}^1$ for which $d_0(F) \geq d_1(F)$. Assume that F has a strong repeller $\mathcal{R}$ that is disjoint from the set of critical values of F. Then, there exist a Borel probability measure μ_F on $\mathbb{P}^1$ that satisfies $F^*(\mu_F) = d_1(F)\mu_F$ and an open set $U(F,\mathcal{R}) \supset \mathcal{R}$ such that*

$$\frac{1}{d_1(F)^n}(F^n)^*(\delta_z) \xrightarrow{weak^*} \mu_F \ \text{ as measures, as } n \to \infty \ \forall z \in U(F,\mathcal{R}). \tag{3.7}$$

In particular, μ_F is invariant under F.

To the best of our knowledge, Theorem 3.4 is the first theorem concerning the equidistribution of preimages of a holomorphic correspondence F such that $d_0(F) \geq d_1(F)$. It would seem that (3.7) could be stronger, since the theorem does not state that $\mathbb{P}^1 \setminus U(F,\mathcal{R})$ is polar (or even nowhere dense). However, given that $d_0(F) \geq d_1(F)$, this is very much in the nature of things. In this regard, we make the following remark.

Remark 3.5. If F is as in Theorem 3.4, we *cannot conclude, in general, that the set $\mathbb{P}^1 \setminus U(F,\mathcal{R})$ is polar.* The following example constitutes a basic obstacle to $\mathbb{P}^1 \setminus U(F,\mathcal{R})$ being even nowhere dense. Let P be any polynomial whose filled Julia set has non-empty interior. Consider the holomorphic correspondence F determined by

$$\Gamma^P \ := \ \text{the completion in } \mathbb{P}^1 \times \mathbb{P}^1 \text{ of the zero set of } (z_1 - P(z_2)).$$

Here, $d_0(F) = \deg(P) > 1 = d_1(F)$. Note that $\{\infty\}$ is a strong repeller. However, $U(F, \{\infty\})$, cannot contain any points from the filled Julia set of P.

The proof of Theorem 3.4 relies on techniques developed by Lyubich in [14]. Given our hypothesis on the existence of a repeller $\mathcal{R}$, one can show that there exists a compact set B such that $\mathcal{R} \subset B^\circ$ and ${}^\dagger F(B) \subset B$. This allows us to define a Perron–Frobenius-type operator $\mathbb{A}_B : \mathcal{C}(B; \mathbb{C}) \longrightarrow \mathcal{C}(B; \mathbb{C})$, where

$$\mathbb{A}_B \ := \ \frac{1}{d_1(F)} \, \boldsymbol{\Lambda}|_B \,,$$

with $\boldsymbol{\Lambda}|_B$ being the operator given by (3.3) with B replacing Ω. Our proof relies on showing that the family $\{\mathbb{A}_B^n : n = 1, 2, 3, \dots\}$ satisfies the conditions of the main result in [14, Section 2]. This goal is achieved, *in part*, by showing that for each $z \in U(F, \mathcal{R})$ there are, for each $n \in \mathbb{Z}_+$, sufficiently many holomorphic branches of n-fold iteration of the correspondence ${}^\dagger F$.

One might ask whether there are lots of holomorphic correspondences F on $\mathbb{P}^1$ with $d_0(F) \geq d_1(F)$ — and *not* just those described in Remark 3.5 — that satisfy the conditions stated in Theorem 3.4. The reader is referred to a short section on constructing examples at the end of this paper. In the next section, we shall establish a few technical facts which will be of relevance throughout this paper. The proofs of our theorems will be provided in Sections 5 and 6.

4. Some basic technical propositions

We begin by showing that the composition of two holomorphic correspondences on $\mathbb{P}^1$, under the composition rule (2.3), produces a holomorphic correspondence.

One way to see this is to begin with how one actually computes $F_2 \circ F_1$ if one is given exact expressions for F_1 and F_2. Let $\mathit{\Gamma}^s$ be the graph of F_s, $s = 1, 2$, and consider the representations given by (2.1). Fix indices j and l such that $1 \leq j \leq L_1$ and $1 \leq l \leq L_2$. It follows that there exist irreducible polynomials $P_1, P_2 \in \mathbb{C}[z, w]$ such that

$$\Gamma^\bullet_{1,j} \cap \mathbb{C}^2 = \{(z, w) \in \mathbb{C}^2 : P_1(z, w) = 0\}, \quad \Gamma^\bullet_{2,l} \cap \mathbb{C}^2 = \{(z, w) \in \mathbb{C}^2 : P_2(z, w) = 0\};$$

see, for instance, [17, pp. 23-24]. Now, given any polynomial $P \in \mathbb{C}[z, w]$, set

$$\begin{aligned} \operatorname{supp}(P) &:= \{(\alpha, \beta) \in \mathbb{N}^2 : \partial_z^\alpha \partial_w^\beta P(0, 0) \neq 0\}, \\ d_z(P) &:= \max\{\alpha \in \mathbb{N} : (\alpha, \beta) \in \operatorname{supp}(P) \text{ for some } \beta \in \mathbb{N}\}, \\ d_w(P) &:= \max\{\beta \in \mathbb{N} : (\alpha, \beta) \in \operatorname{supp}(P) \text{ for some } \alpha \in \mathbb{N}\}. \end{aligned}$$

Then, there is a choice of projective coordinates on $\mathbb{P}^1$ such that

$$\begin{aligned} \Gamma^\bullet_{1,j} &= \{([z_0 : z_1], [w_0 : w_1]) \in \mathbb{P}^1 \times \mathbb{P}^1 : z_0^{d_z(P_1)} w_0^{d_w(P_1)} P_1(z_1/z_0, w_1/w_0) = 0\}, \\ \Gamma^\bullet_{2,l} &= \{([z_0 : z_1], [w_0 : w_1]) \in \mathbb{P}^1 \times \mathbb{P}^1 : z_0^{d_z(P_2)} w_0^{d_w(P_2)} P_2(z_1/z_0, w_1/w_0) = 0\}. \end{aligned}$$

With these notations, we are in a position to state our first proposition.

Proposition 4.1. *Let* $\Gamma^\bullet_{1,j}$ *and* $\Gamma^\bullet_{2,l}$ *be irreducible subvarieties belonging to the holomorphic* 1*-chains* $\mathit{\Gamma}^1$ *and* $\mathit{\Gamma}^2$ *respectively. Let* P_1 *and* P_2 *be the defining functions of* $\Gamma^\bullet_{1,j} \cap \mathbb{C}^2$ *and* $\Gamma^\bullet_{2,l} \cap \mathbb{C}^2$ *respectively.*

- *i) Let* $R(z,w) := \mathsf{Res}(P_1(z,\cdot),P_2(\cdot,w))$*, where* Res *denotes the resultant of two univariate polynomials. Let* V_R *denote the biprojective completion in* $\mathbb{P}^1\times\mathbb{P}^1$ *of* $\{(z,w)\in\mathbb{C}^2 : R(z,w)=0\}$*. Then* $|\Gamma^\bullet_{2,l}\circ\Gamma^\bullet_{1,j}| = V_R$*.*
- *ii)* V_R *has no irreducible components of the form* $\{a\}\times\mathbb{P}^1$ *or* $\mathbb{P}^1\times\{a\}$*,* $a\in\mathbb{P}^1$*.*

Proof. Let us write $V := |\Gamma^\bullet_{2,l}\circ\Gamma^\bullet_{1,j}|$. Since two polynomials $p,q\in\mathbb{C}[X]$ have a common zero if and only if $\mathsf{Res}(p,q)=0$,

$$V\cap\mathbb{C}^2 \ = \ \{(z,w)\in\mathbb{C}^2 : \mathsf{Res}(P_1(z,\cdot),P_2(\cdot,w))=0\}.$$

Hence, as V is the biprojective completion of $V\cap\mathbb{C}^2$ in $\mathbb{P}^1\times\mathbb{P}^1$, (i) follows.

To prove (ii), let us first consider the case when $a\neq[0:1]$. Then, it suffices to show that R has no factors of the form $(z-a)$ or $(w-a)$. We shall show that R has no factors of the form $(z-a)$. A completely analogous argument will rule out factors of the form $(w-a)$, $a\in\mathbb{C}$. To this end, assume that there exists an $a\in\mathbb{C}$ such that $(z-a)|R$ in $\mathbb{C}[z,w]$. This implies

$$R(a,w) \ = \ 0 \ \ \forall w\in\mathbb{C}.$$

Thus, for each $w\in\mathbb{C}$, the polynomial $P_2(\cdot,w)$ has a zero in common with $p_1 := P_1(a,\cdot)\in\mathbb{C}[X]$. Note that $p_1\not\equiv 0$ because, otherwise, $(z-a)|P_1$, which would contradict the fact that $\pi_s|_{\Gamma^\bullet_{1,j}}$ is surjective, $s=1,2$. Thus, there exists an uncountable set $S\subset\mathbb{C}$ and a point $b\in p_1^{-1}\{0\}$ such that

$$P_2(b,w) \ = \ 0 \ \ \forall w\in S.$$

But this implies $P_2(b,\cdot)\equiv 0$, i.e. that $(z-b)|P_2$. This is impossible, for exactly the same reason that $(z-a)\nmid P_1$. Hence R has no factors of the form $(z-a)$, $a\in\mathbb{C}$.

Note that, if we write $\mathbb{C}' := \{[z_0:z_1]\in\mathbb{P}^1 : z_1\neq 0\}$, then, arguing as in the beginning of this proof,

$$V\cap(\mathbb{C}'\times\mathbb{C}) \ = \ \{(z,w)\in\mathbb{C}^2 : z^{d_z(R)}\mathsf{Res}(P_1(1/z,\cdot),P_2(\cdot,w))=0\},$$

where $d_z(R)$ is as defined in the beginning of this section. If we define $\widetilde{R}\in\mathbb{C}[z,w]$ by

$$\widetilde{R}(z,w) \ := \ z^{d_z(R)}\mathsf{Res}(P_1(1/z,\cdot),P_2(\cdot,w)),$$

then an argument analogous to the one above shows that $z\nmid\widetilde{R}$ in $\mathbb{C}[z,w]$. This establishes that $\{[0:1]\}\times\mathbb{P}^1$ is not an irreducible component of V. A similar argument shows that $\mathbb{P}^1\times\{[0:1]\}$ is not an irreducible component of V either. □

Corollary 4.2. *Let* Γ^1 *and* Γ^2 *be two holomorphic correspondences on* $\mathbb{P}^1$*. Then* $\Gamma^2\circ\Gamma^1$ *is a holomorphic correspondence on* $\mathbb{P}^1$*.*

Proof. Let us pick $\Gamma^\bullet_{1,j}$ and $\Gamma^\bullet_{2,l}$ as in Proposition 4.1 and let C be an irreducible component of the variety $|\Gamma^\bullet_{2,l}\circ\Gamma^\bullet_{1,j}|$. We must show that $\pi_s|_C$ is surjective, $s=1,2$. Recalling the explanations given in the proof of Proposition 4.1, it suffices to show the following:

$(**)$ *Given any* $a\in\mathbb{C}$*,* $\pi_s^{-1}\{a\}\cap C\neq\varnothing$*,* $s=1,2$*.*

We shall prove $(**)$ for $s=1$; the proof for $s=2$ follows *mutatis mutandis.* There exists a polynomial $P\in\mathbb{C}[z,w]$ such that $C\cap\mathbb{C}^2=\{(z,w)\in\mathbb{C}^2 : P(z,w)=0\}$ — see, for instance, [17, pp. 23-24]. We may write

$$P(z,w) \ = \sum_{n=0}^{d_w(P)} p_n(z)w^n,$$

where $p_n \in \mathbb{C}[z],\ n = 0,\dots,d_w(P)$, and $d_w(P)$ is as defined at the beginning of this section. It follows from the fundamental theorem of algebra that there exists a point $b \in \mathbb{C}$ such that $P(a,b) = 0$, unless

$$p_0(a) \neq 0 \ \text{ and } \ p_n(a) = 0 \ \forall n \neq 0.$$

This leads to two cases.

Case 1. *The degree* $d_w(P) = 0$.
In this case P depends only on z, in which case C is of the form $\{b\} \times \mathbb{P}^1,\ b \in \mathbb{P}^1$. But, by Part (ii) of Proposition 4.1, this is impossible. Hence, this case cannot arise.

Case 2. *The degree* $d_w(P) > 0$.
In this case, note that

$$z_0^{d_z(P)} w_0^{d_w(P)} P(a/z_0, w_1/w_0)\Big|_{z_0=1,\, w_0=0} \ = \ 0.$$

This means that $([1:a],[0:1]) \in \pi_1^{-1}\{a\} \cap C$.
These two cases together establish $(**)$ for $s = 1$. $\square$

The next lemma will be useful in simplifying expressions of the form $(F^n)^*\delta_z$ or $(F^n)_*\delta_z$.

Lemma 4.3. *Let X be a compact complex manifold and let F be a* holomorphic *correspondence on X. Then* $^\dagger(F^n) = (^\dagger F)^n \ \forall n \in \mathbb{Z}_+$.

Proof. Let $k := \dim(X)$ and let Γ be the holomorphic k-chain that determines F. We shall use the representation

$$\Gamma \ = \ {\sum}'_{j\in[1\,..\,L]} \Gamma^\bullet_j$$

for Γ. We must show that

$$^\dagger(\Gamma^{\circ n}) = (^\dagger\Gamma)^{\circ n} \ \ \forall n \in \mathbb{Z}_+. \tag{4.1}$$

This is clearly true for $n = 1$. Let us assume that (4.1) is true for $n = m$. By this assumption

$$(^\dagger\Gamma)^{\circ(m+1)} \ = \ ^\dagger(\Gamma^{\circ m}) \circ {}^\dagger\Gamma. \tag{4.2}$$

We adopt notation analogous to that used prior to Definition 2.1 For any multi-index $\boldsymbol{\alpha} \in [1\,..\,L]^j$, $\boldsymbol{\alpha} = (\alpha_1,\dots,\alpha_j)$, $j = 1,2,3,\dots$, let us define

$$\begin{aligned} \Gamma^{\bullet\boldsymbol{\alpha}} \ &:= \ \Gamma^\bullet_{\alpha_j} \circ \Gamma^{\bullet(\alpha_1,\dots,\alpha_{j-1})} && \text{if } j \geq 2,\\ \Gamma^{\bullet\boldsymbol{\alpha}} \ &:= \ \Gamma^\bullet_{\alpha_1} && \text{if } j = 1. \end{aligned}$$

Now, for $j = m \geq 2$, where m is as in (4.2), define

$$I(\boldsymbol{\alpha}) \ := \ \text{the collection of all irreducible subvarieties comprising the 1-chain } \Gamma^{\bullet\boldsymbol{\alpha}}$$

(now $\boldsymbol{\alpha} \in [1\,..\,L]^m$). Then, by the composition rules (2.2) and (2.3)

$$^{\dagger}(\Gamma^{\bullet\boldsymbol{\alpha}}) \circ {}^{\dagger}\Gamma^{\bullet}_j \;=\; \sum_{Y\in I(\boldsymbol{\alpha})} \sum_{l=1}^{N(Y,l)} \nu_{j,\,Y,\,l}\, c(Y) Z_{Y,\,l},$$

where the $Z_{Y,\,l}$'s are the distinct irreducible subvarieties of

$$\{(z_1, z_2) \in (\mathbb{P}^1)^2 : \exists w \in \mathbb{P}^1 \text{ s.t. } (w, z_1) \in \Gamma^{\bullet}_j,\ (z_2, w) \in Y\}, \tag{4.3}$$

$c(Y) \in \mathbb{Z}_+$ is the coefficient of Y in the expansion of $\Gamma^{\bullet\boldsymbol{\alpha}}$ according to the representation (1.2), and $\nu_{j,\,Y,\,l} \in \mathbb{Z}_+$ is the generic number w's — as (z_1, z_2) varies through $Z_{Y,\,l}$ — for which the membership conditions in (4.3) are satisfied. From the form of the sets in (4.3), clearly

$$^{\dagger}(\Gamma^{\bullet\boldsymbol{\alpha}}) \circ {}^{\dagger}\Gamma^{\bullet}_j \;=\; {}^{\dagger}(\Gamma^{\bullet}_j \circ \Gamma^{\bullet\boldsymbol{\alpha}}).$$

Combining this with (4.2), we get

$$\begin{aligned}({}^{\dagger}\mathit{\Gamma})^{\circ(m+1)} \;&=\; {\sum}'_{j\in[1\,..\,L]} {\sum}'_{\boldsymbol{\alpha}\in[1\,..\,L]^m} {}^{\dagger}(\Gamma^{\bullet}_j \circ \Gamma^{\bullet\boldsymbol{\alpha}}) \\ &= {}^{\dagger}(\mathit{\Gamma} \circ \mathit{\Gamma}^{\circ m}).\end{aligned}$$

The expression (4.1) follows by induction. □

The final result in this section is important because it establishes that the measures $d_1(F)^{-n}(F^n)^*(\delta_z)$, occurring in Theorems 3.1 and 3.4, are probability measures. *The result below is obvious if F is a map.* Its proof is simple but, for general correspondences, a little care is needed to account for multiplicities.

Proposition 4.4. *Let F be a holomorphic correspondence on $\mathbb{P}^1$. Then $d_{top}(F^n) = d_{top}(F)^n\ \forall n \in \mathbb{Z}_+$.*

Remark 4.5. As a matter of fact, Proposition 4.4 is true for holomorphic correspondences on *any* compact complex manifold X, and not just on $\mathbb{P}^1$. The proof is almost exactly the one below, except that when $X \neq \mathbb{P}^1$ those parts of the proof below that rely on Part (*ii*) of Proposition 4.1 will, in the general case, follow from the following:

> *Let X be a compact complex manifold and let $\Gamma^{\bullet}_{1,\,j}$ and $\Gamma^{\bullet}_{2,\,l}$ be as in* (2.1). *The projections π_1 and π_2 are surjective when restricted to each irreducible component of $|\Gamma^{\bullet}_{2,\,l} \circ \Gamma^{\bullet}_{1,\,j}|$.*

Since we have not given a proof of the above when $X \neq \mathbb{P}^1$ — doing so would be a considerable digression — we state Proposition 4.4 for $X = \mathbb{P}^1$ only.

Proof. We shall use induction. The above formula is a tautology for $n = 1$. Assume that it is true for $n = m$ for some $m \in \mathbb{Z}_+$. Let $\mathit{\Gamma}$ be the holomorphic 1-chain that determines F. Following the representation (1.2), let us denote

$$\mathit{\Gamma}^{\circ m} \;=\; \sum_{j=1}^{M} \mu_j \boldsymbol{\Lambda}_j,$$

and let $\mathit{\Gamma}$ be precisely as in (1.2). Then

$$\mathit{\Gamma}^{\circ(m+1)} \ = \ \sum_{j=1}^{M}\sum_{l=1}^{N}\mu_j m_l(\Gamma_l\circ\boldsymbol{\Lambda}_j).$$

Let $\widetilde{V}$ be a Zariski-open subset of $\mathbb{P}^1$ such that $(\pi_2^{-1}(\widetilde{V})\cap\Gamma_l,\widetilde{V},\pi_2)$, $(\pi_2^{-1}(\widetilde{V})\cap\boldsymbol{\Lambda}_j,\widetilde{V},\pi_2)$ and $(\pi_1^{-1}(\widetilde{V})\cap\boldsymbol{\Lambda}_j,\widetilde{V},\pi_1)$ are covering spaces for each $(j,l)\in[1\,..\,M]\times[1\,..\,N]$. We can then find a Zariski-open subset V of $(\mathbb{P}^1)^2$ such that

($\bullet$) $V\subset\widetilde{V}^2$, $(\widetilde{V}^2\setminus V)$ is an at most finite union of lines of the form $\{a\}\times\mathbb{P}^1$ or $\mathbb{P}^1\times\{a\}$, and for each $(j,l)\in[1\,..\,M]\times[1\,..\,N]$, $\pi_2^{-1}\{w\}\cap V\supset\pi_2^{-1}\{w\}\cap\Gamma_l$ and $\pi_1^{-1}\{z\}\cap V\supset\pi_1^{-1}\{z\}\cap\boldsymbol{\Lambda}_j$ $\forall(z,w)\in V$.

That there is a V for which ($\bullet$) holds true for all pertinent (j,l) follows easily from Part *(ii)* of Proposition 4.1. (We shall provide an argument below for a fact analogous to ($\bullet$); a simpler version that argument gives ($\bullet$).)

Let us write

$$\Delta(\boldsymbol{\Lambda},j) \ := \ \text{the number of sheets of } (\pi_2^{-1}(\pi_2(V))\cap\boldsymbol{\Lambda}_j,\pi_2(V),\pi_2),$$
$$\Delta(\mathit{\Gamma},l) \ := \ \text{the number of sheets of } (\pi_2^{-1}(\pi_2(V))\cap\Gamma_l,\pi_2(V),\pi_2).$$

We now fix a $(j,l)\in[1\,..\,M]\times[1\,..\,N]$. Let us write $X_{jl}:=|\Gamma_l\circ\boldsymbol{\Lambda}_j|$. Let $Y_{s,\,jl}$'s, $s=1,\dots,N(j,l)$, be the distinct irreducible components of X_{jl}. It follows from a standard intersection-theory argument (note that $Y_{s,\,jl}$ is *irreducible*), the fact that $\dim_{\mathbb{C}}(Y_{s,\,jl})=1$, and from the structure of V, that there is subset $W\subset\mathbb{P}^1$ such that W^2 is a Zariski-open subset of V and

$$\mathsf{Card}\left(\pi_1(\pi_2^{-1}\{w\}\cap\Gamma_l)\cap\pi_2(\pi_1^{-1}\{z\}\cap\boldsymbol{\Lambda}_j)\right) \ = \ \nu_{s,\,jl} \ \ \forall(z,w)\in Y_{s,\,jl}\cap W^2, \tag{4.4}$$

for each $s=1,\dots,N(j,l)$ and $(j,l)\in[1\,..\,M]\times[1\,..\,N]$, where $\nu_{s,\,jl}$ is as given immediately after the composition rule (2.2).

Consider the set

$$\Omega_{jl} \ := \ \{(z,w)\in W^2 : \pi_2^{-1}\{w\}\cap W^2\supset\pi_2^{-1}\{w\}\cap\Gamma_l \text{ and } \pi_1^{-1}\{z\}\cap W^2\supset\pi_1^{-1}\{z\}\cap\boldsymbol{\Lambda}_j\}.$$

This set is clearly a Zariski-open subset of W^2, but we claim that Ω_{jl} is a *non-empty* Zariski-open set. To this end, consider the sets

$$E_1 \ := \ (V\setminus W^2)\cap\boldsymbol{\Lambda}_j, \qquad E_2 \ := \ (V\setminus W^2)\cap\Gamma_l.$$

Owing to the Part *(ii)* of Proposition 4.1 and the structure of V, E_1 and E_2 are finite sets. Hence

$$W^2\setminus\left(\pi_2^{-1}(\pi_2(E_2))\cup\pi_1^{-1}(\pi_1(E_1))\right) \ \neq \ \varnothing.$$

Then, owing to the property ($\bullet$)

$$\Omega_{jl}\supset W^2\setminus\left(\pi_2^{-1}(\pi_2(E_2))\cup\pi_1^{-1}(\pi_1(E_1))\right),$$

whence our claim. Now, for each $w\in\pi_2(\Omega_{jl})$, define

$$S_{jl}(w) \ := \bigcup_{(\zeta,w)\in\pi_2^{-1}\{w\}\cap\Gamma_l}\pi_2^{-1}\{\zeta\}\cap\boldsymbol{\Lambda}_j.$$

By the construction of Ω_{jl}, $\pi_2^{-1}\{w\} \cap \Gamma_l \subset W^2 \subset V \ \forall w \in \pi_2(\Omega_{jl})$. Then, by the construction of V,

$$\mathsf{Card}(S_{jl}(w)) \ = \ \Delta(\mathbf{\Lambda}, j)\Delta(\mathit{\Gamma}, l) \quad \forall w \in \pi_2(\Omega_{jl}). \tag{4.5}$$

Let $\mathfrak{S}_{jl}$ denote the set of all points in X_{jl} that belong to more than one $Y_{s,\,jl}$. Then, by (4.5)

$$\sum_{s=1}^{N(j,l)} \nu_{s,\,jl}\mathsf{Card}(\pi_2^{-1}\{w\} \cap Y_{s,\,jl}) \ = \ \Delta(\mathbf{\Lambda}, j)\Delta(\mathit{\Gamma}, l) \quad \forall w \in \pi_2(\Omega_{jl}) \setminus \pi_2(\mathfrak{S}_{jl}). \tag{4.6}$$

Recalling our definitions:

$$d_{top}(F) \ = \ \sum_{l=1}^{N} m_l\, \Delta(\mathit{\Gamma}, l), \qquad d_{top}(F^m) \ = \ \sum_{j=1}^{M} \mu_j\, \Delta(\mathbf{\Lambda}, j). \tag{4.7}$$

Now let V' be a Zariski-open subset of $\mathbb{P}^1$ such that $(\pi_2^{-1}(V') \cap Y_{s,\,jl}, V', \pi_2)$ are covering spaces for every $s = 1, \dots, N(j,l)$ and every $(j,l) \in [1\,..\,M] \times [1\,..\,N]$. Then, writing $\Omega := (V' \cap \pi_2\,(\cap_{j,\,l}\Omega_{jl})) \setminus \pi_2\,(\cup_{j,\,l}\mathfrak{S}_{jl})$, we have by definition

$$d_{top}(F^{m+1}) \ = \ \sum_{j=1}^{M}\sum_{l=1}^{N}\sum_{s=1}^{N(j,l)} \mu_j\, m_l\, \nu_{s,\,jl}\, \mathsf{Card}(\pi_2^{-1}\{w\} \cap Y_{s,\,jl}) \quad \text{for any } w \in \Omega.$$

In view of (4.6) and (4.7), the above equation simplifies as follows:

$$\begin{aligned} d_{top}(F^{m+1}) \ &= \ \sum_{j=1}^{M}\sum_{l=1}^{N} \mu_j\, m_l\, \Delta(\mathbf{\Lambda}, j)\Delta(\mathit{\Gamma}, l) \\ &= \ d_{top}(F^m)\, d_{top}(F) \\ &= \ d_{top}(F)^m\, d_{top}(F) \qquad\qquad \text{[by the inductive hypothesis].} \end{aligned}$$

By induction, the result follows. $\square$

5. The proof of Theorem 3.1

As stated earlier, Part (i) of Theorem 3.1 follows almost directly from some of the results in [9]. We therefore begin this section with a statement of the results needed. To do so, we need a definition. If (X, ω) is a compact Kähler manifold of dimension k, ω so normalised that $\int_X \omega^k = 1$, then for each $s = 0, 1, \dots, k$, the number

$$\lambda_s(F) \ := \ \int_X F^*(\omega^s) \wedge \omega^{k-s}$$

is called the *intermediate degree of F of order s*. In view of the prescription (3.1), the above definition can be rewritten as

$$\lambda_s(F) \ = \ \sum_{j=1}^{N} m_j \int_{\mathrm{reg}(\Gamma_j)} \Big(\pi_1|_{\mathrm{reg}(\Gamma_j)}\Big)^* \omega^{k-s} \wedge \Big(\pi_2|_{\mathrm{reg}(\Gamma_j)}\Big)^* \omega^s, \tag{5.1}$$

with the understanding that $\omega^0 := 1$, and that $\operatorname{reg}(\Gamma_j)$ is the set of all points at which Γ_j is smooth. For $s = 0, 1, \dots, k$, we define the *dynamical degree of F of order s* as

$$d_s(F) \ := \ \limsup_{n\to\infty}(\lambda_s(F^n))^{1/n}.$$

We now state two essential results. In this section, whenever we refer to a compact Kähler manifold (X, ω), we will assume that ω is normalised so that $\int_X \omega^k = 1$.

Result 5.1 (Theorem 4.6 of [9] paraphrased for correspondences). *Let (X, ω) be a compact Kähler manifold of dimension k and let $F_n : X \longrightarrow X$, $n = 1, 2, 3, \dots$, be meromorphic correspondences on X. Suppose the series $\sum_{n\in\mathbb{Z}_+} d_{top}(F_n)^{-1}\lambda_{k-1}(F_n)$ converges. Then, there exists a pluripolar subset $\mathcal{E}$ of X such that for each $x \in X \setminus \mathcal{E}$,*

$$\left\langle d_{top}(F_n)^{-1}\left(F_n^*(\delta_x) - F_n^*(\omega^k)\right), \psi\right\rangle \longrightarrow 0 \ \text{ as } n \to \infty$$

for each $\psi \in \mathcal{C}^2(X;\mathbb{R})$ and, in fact, this convergence is uniform on subsets that are bounded with respect to the $\mathcal{C}^2$-norm.

Result 5.2 (a version of Corollary 5.3, [9]). *Let F be a meromorphic correspondence on a compact Kähler manifold (X, ω) of dimension k. Suppose $d_{k-1}(F) < d_{top}(F)$. Then, there exists a PLB probability measure μ_F such that*

$$\frac{1}{d_{top}(F)^n}(F^n)^*(\omega^k) \xrightarrow{\text{weak*}} \mu_F \ \text{ as measures, as } n \to \infty.$$

Furthermore, μ_F satisfies $F^*\mu_F = d_{top}(F)\mu_F$.

We refer the reader to [9, Section 2.2] for a definition of PLB measures; what is of relevance to us is that μ_F is a Borel probability measure.

The reader will note that one quantitative departure from the original in our paraphrasing of [9, Theorem 4.6] is that we have replaced all occurrences of $\lambda_k(F_n)$ in the original by $d_{top}(F_n)$. This is because, in the case of meromorphic *correspondences*, $\lambda_k(F_n) = d_{top}(F_n)$. Theorem 4.6 of [9] applies to the more general class of multivalued maps that Dinh–Sibony call *meromorphic transforms*, for which the last equality may not hold true. Before proceeding further, we justify this equality.

Lemma 5.3. *Let (X, ω) be a compact Kähler manifold of dimension k and let $F : X \longrightarrow X$ be a meromorphic correspondence. Then, $\lambda_k(F) = d_{top}(F)$.*

Proof. Let Γ be the graph of F, and consider the representation (1.2) for it. Let C denote an irreducible component occurring in the representation (1.2). Let Ω be a Zariski-open subset of X such that $(\pi_2^{-1}(\Omega)\cap C, \Omega, \pi_2)$ is a covering and $E := \pi_2^{-1}(\Omega)\cap C$ is a smooth manifold. Let Δ be the degree of this covering.

Let us write

$$I(C, \omega) \ := \ \int_{\operatorname{reg}(C)} \left(\pi_2|_{\operatorname{reg}(C)}\right)^* \omega^k.$$

By the definition of $\operatorname{reg}(C)$, $\left(\pi_2|_{\operatorname{reg}(C)}\right)^*\omega^k$ vanishes at each point on $\operatorname{reg}(C) \setminus E$. Thus,

$$I(C, \omega) \ = \ \int_E \left(\pi_2|_{\operatorname{reg}(C)}\right)^* \omega^k. \tag{5.2}$$

As $(\pi_2|_E)^*\omega^k$ is a volume form, we have an associated Lebesgue measure on E. Then, the integral in (5.2) is the Lebesgue integral of a certain non-negative continuous function $f_{C,\omega} : E \longrightarrow [0, \infty)$. Furthermore, observe that:

- $\pi_2|_E$ is Lipschitz relative to any smooth local coordinates on it.
- As $I(C, \omega)$ is finite, $f_{C,\omega}$ is integrable with respect to the Lebesgue measure on E.

The last three facts provide us the setting in which to apply the change-of-variable formula for the map $\pi_2|_E$, which gives us

$$\int_E \Big(\pi_2|_{\text{reg}(C)}\Big)^* \omega^k \ = \ \Delta \int_\Omega \omega^k.$$

From this, (5.2), and the fact that Ω is Zariski-open in X, it follows that

$$I(C, \omega) \ = \ \Delta \int_\Omega \omega^k \ = \ \Delta \tag{5.3}$$

(owing to the normalisation of ω). Now let $\Delta(\mathit{\Gamma}, j)$ denote the cardinality of $\pi_2^{-1}\{x\} \cap \Gamma_j$ for generic x. Since (5.3) holds true for any irreducible component C that constitutes $\mathit{\Gamma}$, it follows that

$$\lambda_k(F) \ = \ \sum_{j=1}^N m_j \Delta(\mathit{\Gamma}, j) \ = \ d_{top}(F).$$

$\square$

Note: the above lemma also establishes that the numbers $d_0(F)$ and $d_1(F)$ occurring in the statement of Theorem 3.1 agree with the dynamical degrees of order 0 and 1 respectively.

The proof of Theorem 3.1. i) With $\omega = \omega_{FS}$, it follows from Lemma 5.3 and the definition of $\lambda_0(F)$ that $\lambda_0(F) = d_{top}({}^{\dagger}F) = d_0(F)$. Therefore, in view of Proposition 4.4, the series

$$\sum\nolimits_{n \in \mathbb{Z}_+} d_{top}(F^n)^{-1} \lambda_{k-1}(F^n) \ = \ \sum\nolimits_{n \in \mathbb{Z}_+} d_1(F)^{-n} d_0(F)^n \quad \text{is convergent.}$$

Therefore, from Result 5.1, it follows that there is a polar set $\mathcal{E} \subsetneq \mathbb{P}^1$ such that for each $z \in \mathbb{P}^1 \setminus \mathcal{E}$

$$\big\langle d_{top}(F)^{-n} \left((F^n)^*(\delta_z) - (F^n)^*(\omega_{FS}) \right), \psi \big\rangle \longrightarrow 0 \ \text{ as } n \to \infty, \tag{5.4}$$

where $\psi \in \mathcal{C}^2(\mathbb{P}^1; \mathbb{R})$. By hypothesis, Result 5.2 is applicable. Let μ_F be the probability measure guaranteed by Result 5.2. From the discussion culminating in (3.4) we see that, for each $n \in \mathbb{Z}_+$, $(F^n)^*(\delta_z)$ *can be viewed as a measure.* Hence, combining the conclusion of Result 5.2 with (5.4), we see that if $z \in \mathbb{P}^1 \setminus \mathcal{E}$, then

$$d_{top}(F)^{-n} \int_{\mathbb{P}^1} \psi \, d(F^n)^*(\delta_z) \longrightarrow \int_{\mathbb{P}^1} \psi \, d\mu_F \ \text{ as } n \to \infty \tag{5.5}$$

for each fixed $\psi \in \mathcal{C}^2(\mathbb{P}^1; \mathbb{R})$.

Let us fix a $z \in \mathbb{P}^1 \setminus \mathcal{E}$. It follows from (3.4) and Proposition 4.4 that $d_{top}(F)^{-n}(F^n)^*(\delta_z)$ is a probability measure. With this fact, it is easy to see that the mode of convergence in (5.5) implies the conclusion of Part (i). To see this, let us write $d_{top}(F)^{-n}(F^n)^*(\delta_z)$

as ν_n^z. Now fix a *continuous* function $f \in \mathcal{C}(\mathbb{P}^1;\mathbb{R})$. Given an $\varepsilon > 0$, we may pick a $\psi \in \mathcal{C}^\infty(\mathbb{P}^1;\mathbb{R})$ such that $\sup_{\mathbb{P}^1} |f - \psi| < \varepsilon/3$. Applying (5.5) to *this particular* ψ, and using the triangle inequality, we can find an $N = N(\varepsilon) \in \mathbb{Z}_+$ such that:

$$\left| \int_{\mathbb{P}^1} f \, d\nu_n^z - \int_{\mathbb{P}^1} f \, d\mu_F \right| \; < \; \varepsilon \quad \forall n \geq N.$$

So, for each fixed $f \in \mathcal{C}(\mathbb{P}^1;\mathbb{R})$, $\int_{\mathbb{P}^1} f \, d\nu_n^z \longrightarrow \int_{\mathbb{P}^1} f \, d\mu_F$ as as $n \to \infty$. This is precisely the desired conclusion.

As μ_F is a Borel measure, it is a regular measure. Thus, keeping in mind the approximation trick above, it suffices to show that for any open set U and any continuous function $\varphi : \mathbb{P}^1 \longrightarrow [0,1]$ with $\text{supp}(\varphi) \subset U$, $\int_{\mathbb{P}^1} \mathbf{\Lambda}[\varphi] \, d\mu_F = \int_{\mathbb{P}^1} \varphi \, d\mu_F$, where $\mathbf{\Lambda}[\varphi]$ is as explained in the passage following (3.3). Fix a point $z \notin \mathcal{E}$. By the equidistribution result just established:

$$\begin{aligned} \int_{\mathbb{P}^1} \mathbf{\Lambda}[\varphi] \, d\mu_F \; &= \; \lim_{n\to\infty} d_{top}(F)^{-(n+1)} \left\langle (F^n)^*(\delta_z), \mathbf{\Lambda}[\varphi] \right\rangle \\ &= \; \lim_{n\to\infty} d_{top}(F)^{-(n+1)} \left\langle (F^{n+1})^*(\delta_z), \varphi \right\rangle \; = \; \int_{\mathbb{P}^1} \varphi \, d\mu_F. \quad \text{[by (3.4) above]} \end{aligned}$$

By the preceding discussion, this establishes that μ_F is invariant under F.

(ii) We assume that $\mathcal{N}(F) \neq \varnothing$; there is nothing to prove otherwise. Let us fix a $z_0 \in \mathcal{N}(F)$. Then, any $\boldsymbol{\mathcal{Z}} \in \mathscr{P}_n(F, z_0)$ is admissible, and by Property (4) of Definition 2.1, assuming that $n \geq 2$, we get

$$\begin{aligned} \pi_{j-1}^0 \circ \nu_{(\boldsymbol{\mathcal{Z}}, j-1)} \circ \psi_{(\boldsymbol{\mathcal{Z}}, j-1)} \; &= \pi_{j-1}^0 \circ \mathsf{proj}_j \circ \nu_{(\boldsymbol{\mathcal{Z}}, j)} \circ \psi_{(\boldsymbol{\mathcal{Z}}, j)} \\ &= \pi_j^0 \circ \nu_{(\boldsymbol{\mathcal{Z}}, j)} \circ \psi_{(\boldsymbol{\mathcal{Z}}, j)} \quad \forall j = 2, \dots, n. \end{aligned}$$

Iterating this argument, we deduce the following:

$$\begin{aligned} &\text{For any } n \geq 2, \; \boldsymbol{\mathcal{Z}} \in \mathscr{P}_n(F, z_0) \text{ (and admissible)} \\ &\qquad \Rightarrow \; \pi_j^0 \circ \nu_{(\boldsymbol{\mathcal{Z}}, j)} \circ \psi_{(\boldsymbol{\mathcal{Z}}, j)} \; = \; \pi_1^0 \circ \nu_{(\boldsymbol{\mathcal{Z}}, 1)} \circ \psi_{(\boldsymbol{\mathcal{Z}}, 1)} \quad \forall j = 1, \dots, n. \qquad (5.6) \end{aligned}$$

Let us now *fix* a disc Δ around $0 \in \mathbb{C}$ such that $\Delta \Subset D$ (where D is as given by Definition 2.3). Define

$$K^{z_0} \; := \; \bigcap\nolimits_{j \in [1\,..\,L]} \; \bigcap_{x \in \pi_1^{-1}\{z_0\} \cap \Gamma_j^\bullet} \; \pi_1^0 \circ \nu_{((z_0, \pi_2(x);\, j), 1)} \circ \psi_{((z_0, \pi_2(x);\, j), 1)}(\overline{\Delta}).$$

Clearly, there is a region $G \Subset D$, containing 0, such that

$$\begin{aligned} &(\nu_{x\bullet j} \circ \psi_{x\bullet j})^{-1} \left(\left(\pi_1^0 \big|_{U(x\bullet j)} \right)^{-1} (K^{z_0}) \right) \; \Subset \; G \\ &\qquad\qquad\qquad\qquad \text{for each } x \in \pi_1^{-1}\{z_0\} \cap \Gamma_j^\bullet \text{ and each } j = 1, \dots, L, \end{aligned}$$

where $x \bullet j$ is our abbreviation for the triple $((z_0, \pi_2(x); j), 1)$ ($\equiv ((z_0, \pi_2(x)), j, 1)$ alternatively) and $U(x \bullet j)$ is as described in condition (2) of Definition 2.1. From the above fact and (5.6) we deduce the following:

for any $n \geq 1$ and $\boldsymbol{\mathcal{Z}} = (Z;\, \boldsymbol{\alpha}) \in \mathscr{P}_n(F, z_0)$,

$$\left(\nu_{(\boldsymbol{\mathcal{Z}},n)} \circ \psi_{(\boldsymbol{\mathcal{Z}},n)}\right)^{-1}\left(\left(\pi_n^0\big|_{U(Z,\boldsymbol{\alpha},n)}\right)^{-1}(K^{z_0})\right) \Subset G. \tag{5.7}$$

We can deduce from Definitions 2.1 and 2.3 that $K^{z_0} \subset \mathcal{N}(F)$. As K^{z_0} has non-empty interior, it suffices to show that for any *non-negative* function $\varphi \in \mathcal{C}(\mathbb{P}^1; \mathbb{R})$ with $\operatorname{supp}(\varphi) \subset K^{z_0}$, $\int_{\mathbb{P}^1} \varphi\, d\mu_F = 0$. Hence, let us pick some function $\varphi \in \mathcal{C}(\mathbb{P}^1; \mathbb{R})$ as described. For any path $\boldsymbol{\mathcal{Z}} = (Z, \boldsymbol{\alpha}) \in \mathscr{P}_n(F, z_0)$, let us write

$$\boldsymbol{\mathcal{Z}}^\natural := \left(\pi_n^0 \times \pi_n^n\right) \circ \nu_{(\boldsymbol{\mathcal{Z}},n)} \circ \psi_{(\boldsymbol{\mathcal{Z}},n)}.$$

Then, following the notation introduced in the proof of Lemma 4.3, $\boldsymbol{\mathcal{Z}}^\natural(D)$ is a $\Gamma^{\bullet\boldsymbol{\alpha}}$-open neighbourhood of the point $(\pi_n^0 \times \pi_n^n)(Z)$. Furthermore, by our constructions in Definition 2.1, $\pi_1^{-1}(K^{z_0}) \cap |\varGamma^{\circ n}|$ is covered by the sets $\boldsymbol{\mathcal{Z}}^\natural(D)$ as $\boldsymbol{\mathcal{Z}}$ varies through $\mathscr{P}_n(F, z_0)$. Hence, by definition:

$$\langle (F^n)^*(\omega_{FS}), \varphi \rangle = \sum_{\boldsymbol{\mathcal{Z}} \in \mathscr{P}_n(F,z_0)} \int\limits_{\operatorname{reg}(\boldsymbol{\mathcal{Z}}^\natural(D))} \pi_1^*(\varphi)\pi_2^*(\omega_{FS}).$$

By condition (2) of Definition 2.1, $\nu_{(\boldsymbol{\mathcal{Z}},n)}^{-1}\{p\}$ has cardinality ≥ 2 for $p \in U(Z, \boldsymbol{\alpha}, n)$ if and only if p is a nodal singularity (see Remark 2.2). It is routine to show that $\pi_n^0 \times \pi_n^n$ would map two distinct points $p, q \in U(Z, \boldsymbol{\alpha}, n)$ to the same point precisely when there is at least one j, $1 \leq j \leq n$, such that $\pi_j^j\big|_{U(Z,\boldsymbol{\alpha},j)}$ is not injective. Thus, if $\pi_n^0 \times \pi_n^n\big|_{U(Z,\boldsymbol{\alpha},n)}$ is not injective, then, by condition (2) of Definition 2.1, it has a well-defined topological degree. We shall denote it by $\deg(\boldsymbol{\mathcal{Z}}^\natural)$. By arguments analogous to those in the proof of Lemma 5.3, we can apply the change-of-variables formula to get:

$$\langle (F^n)^*(\omega_{FS}), \varphi \rangle = \sum_{\boldsymbol{\mathcal{Z}} \in \mathscr{P}_n(F,z_0)} \frac{1}{\deg(\boldsymbol{\mathcal{Z}}^\natural)} \int\limits_{\psi_{(\boldsymbol{\mathcal{Z}},n)}(D)} (\varphi \circ \pi_n^0 \circ \nu_{(\boldsymbol{\mathcal{Z}},n)})(\pi_n^n \circ \nu_{(\boldsymbol{\mathcal{Z}},n)})^*(\omega_{FS}).$$

For each $\boldsymbol{\mathcal{Z}} \in \mathscr{P}_n(F, z_0)$, let us write

$$\deg(\boldsymbol{\mathcal{Z}}) := \text{the degree of the map } \psi_{(\boldsymbol{\mathcal{Z}},n)} : D \longrightarrow \nu_{(\boldsymbol{\mathcal{Z}},n)}^{-1}(U(Z, \boldsymbol{\alpha}, n)).$$

We apply the change-of-variables formula for ramified coverings of finite degree to get:

$$\begin{aligned} &\langle (F^n)^*(\omega_{FS}), \varphi \rangle \\ &= \sum_{\boldsymbol{\mathcal{Z}} \in \mathscr{P}_n(F,z_0)} \frac{1}{\deg(\boldsymbol{\mathcal{Z}})\deg(\boldsymbol{\mathcal{Z}}^\natural)} \int\limits_D (\varphi \circ \pi_n^0 \circ \nu_{(\boldsymbol{\mathcal{Z}},n)} \circ \psi_{(\boldsymbol{\mathcal{Z}},n)})(\pi_n^n \circ \nu_{(\boldsymbol{\mathcal{Z}},n)} \circ \psi_{(\boldsymbol{\mathcal{Z}},n)})^*(\omega_{FS}) \\ &= \sum_{\boldsymbol{\mathcal{Z}} \in \mathscr{P}_n(F,z_0)} \frac{1}{\deg(\boldsymbol{\mathcal{Z}})\deg(\boldsymbol{\mathcal{Z}}^\natural)} \int\limits_{\overline{G}} (\varphi \circ \pi_n^0 \circ \nu_{(\boldsymbol{\mathcal{Z}},n)} \circ \psi_{(\boldsymbol{\mathcal{Z}},n)})(\pi_n^n \circ \nu_{(\boldsymbol{\mathcal{Z}},n)} \circ \psi_{(\boldsymbol{\mathcal{Z}},n)})^*(\omega_{FS}) \end{aligned} \tag{5.8}$$

The last expression arises from (5.7) and the fact that $\operatorname{supp}(\varphi) \subset K^{z_0}$.

Endow $\mathbb{P}^1$ with homogeneous coordinates. In view of the argument made below, we may assume without loss of generality that $\pi^n_n \circ \nu_{(\boldsymbol{\mathscr{Z}},n)} \circ \psi_{(\boldsymbol{\mathscr{Z}},n)}(D)$ does not contain *both* $[0:1]$ and $[1:0]$ (if not, then for each n and each $\boldsymbol{\mathscr{Z}} \in \mathscr{P}_n(F, z_0)$ for which this happens, we split the relevant integral below into a sum of integrals over a two-set partition of D). Writing

$$\pi^n_n \circ \nu_{(\boldsymbol{\mathscr{Z}},n)} \circ \psi_{(\boldsymbol{\mathscr{Z}},n)} \;=\; [X_{(\boldsymbol{\mathscr{Z}},n)} : Y_{(\boldsymbol{\mathscr{Z}},n)}],$$

where $X_{(\boldsymbol{\mathscr{Z}},n)}, Y_{(\boldsymbol{\mathscr{Z}},n)} \in \mathcal{O}(D)$ and have no common zeros in D, define

$$q_{(\boldsymbol{\mathscr{Z}},n)} \;:=\; \begin{cases} X_{(\boldsymbol{\mathscr{Z}},n)}/Y_{(\boldsymbol{\mathscr{Z}},n)}, & \text{if } [0:1] \in \pi^n_n \circ \nu_{(\boldsymbol{\mathscr{Z}},n)} \circ \psi_{(\boldsymbol{\mathscr{Z}},n)}(D), \\ Y_{(\boldsymbol{\mathscr{Z}},n)}/X_{(\boldsymbol{\mathscr{Z}},n)}, & \text{if } [1:0] \in \pi^n_n \circ \nu_{(\boldsymbol{\mathscr{Z}},n)} \circ \psi_{(\boldsymbol{\mathscr{Z}},n)}(D). \end{cases}$$

From the expression for the Fubini–Study metric in local coordinates and from (5.8), we have the estimate

$$d_1(F)^{-n} |\langle (F^n)^*(\omega_{FS}), \varphi\rangle| \;\leq\; \frac{C}{d_1(F)^n} \sum_{\boldsymbol{\mathscr{Z}} \in \mathscr{P}_n(F,z_0)} \int\limits_{G} \|\varphi\|_\infty \left(\frac{|q'_{(\boldsymbol{\mathscr{Z}},n)}(\zeta)|}{1 + |q_{(\boldsymbol{\mathscr{Z}},n)}(\zeta)|^2} \right)^2 dA(\zeta).$$

Since, by hypothesis, $\mathscr{F}(z_0)$ is a normal family, it follows — see, for instance, Conway [7, Chapter VII/§3] — that the family

$$\left\{ \frac{|q'_{(\boldsymbol{\mathscr{Z}},n)}|}{1 + |q_{(\boldsymbol{\mathscr{Z}},n)}|^2} : n \in \mathbb{Z}_+ \text{ and } \boldsymbol{\mathscr{Z}} \in \mathscr{P}_n(F, z_0) \right\}$$

is locally uniformly bounded. As $G \Subset D$, there exists an $M > 0$ such that

$$\frac{|q'_{(\boldsymbol{\mathscr{Z}},n)}(\zeta)|}{1 + |q_{(\boldsymbol{\mathscr{Z}},n)}(\zeta)|^2} \;\leq\; M \;\; \forall \zeta \in \overline{G}, \; n \in \mathbb{Z}_+ \text{ and } \forall \boldsymbol{\mathscr{Z}} \in \mathscr{P}_n(F, z_0).$$

Given $n \in \mathbb{Z}_+$, the number of summands in (5.8) is at most $d_{top}({}^{\dagger}(F^n)) = d_0(F)^n$. The latter equality is a consequence of Proposition 4.4 and Lemmas 4.3 and 5.3. From it and from the last two estimates, it follows that

$$d_1(F)^{-n} |\langle (F^n)^*(\omega_{FS}), \varphi\rangle| \;\leq\; C \left[\frac{d_0(F)}{d_1(F)} \right]^n M^2 \mathsf{Area}(\overline{G}) \;\longrightarrow\; 0 \text{ as } n \to \infty, \tag{5.9}$$

since, by hypothesis, $d_0(F) < d_1(F)$.

In view of Result 5.2, with $\omega = \omega_{FS}$, and (5.9), we have

$$\int_{\mathbb{P}^1} \varphi \, d\mu_F \;=\; \lim_{n \to \infty} \frac{1}{d_1(F)^n} \langle (F^n)^*(\omega_{FS}), \varphi\rangle \;= 0.$$

By our remarks earlier, (ii) follows. ☐

6. The proof of Theorem 3.4

As the paragraphs in Section 3 preceding the statement of Theorem 3.4 would suggest, its proof relies on several notions introduced in [15]. We therefore begin this section with a definition and a couple of results from [15].

Definition 6.1. Let X be a compact Hausdorff space and let $f \subset X \times X$ be a relation of X to itself such that $\pi_1(f) = X$. A set $B \subset X$ is called an *attractor block* for f if $f(\overline{B}) \subset B^\circ$.

We recall that, given a relation f and a set $S \subset X$, $f(S)$ is as defined in Section 3.

Result 6.2 (McGehee, Theorem 7.2 of [15])**.** *Let X be a compact Hausdorff space and let $f \subset X \times X$ be a relation of X to itself such that $\pi_1(f) = X$. Assume f is a closed set. If B is an attractor block for f, then B is a neighbourhood of $\omega(B; f)$: i.e., $\omega(B; f) \subset B^\circ$.*

Result 6.3 (McGehee, Theorem 7.3 of [15])**.** *Let X be a compact Hausdorff space and let $f \subset X \times X$ be a relation of X to itself such that $\pi_1(f) = X$. Assume f is a closed set. If $\mathcal{A}$ is an attractor for f and V is a neighbourhood of $\mathcal{A}$, then there exists a closed attractor block B for f such that $B \subset V$ and $\omega(B; f) = \mathcal{A}$.*

Before we can give the proof of Theorem 3.4, we need one more concept. For this purpose, we shall adapt some of the notations developed in Section 2.2. Here, F will denote a holomorphic correspondence on $\mathbb{P}^1$. Firstly: given $N \in \mathbb{Z}_+$, we say that $(w_0, w_{-1} \dots, w_{-N}; \alpha_1, \dots, \alpha_N) \in (\mathbb{P}^1)^{N+1} \times [1\,..\,L]^N$ (see (2.1) for the meaning of L) is *a path of a backward iteration of F starting at w_0, of length N*, if

$$(w_{-j}, w_{-j+1}) \in \Gamma^\bullet_{\alpha_j}, \quad j = 1, \dots, N.$$

In analogy with the notation in Section 2.2, we set:

$$\mathscr{P}_{-N}(F, w_0) \; := \; \text{the set of all paths of } \textit{backward} \text{ iterations} \\ \text{of } F, \text{ of length } N, \text{ starting at } w_0.$$

Next, we say that a point $w \in \mathbb{P}^1$ is a *regular value* of F if it is not a critical value (recall that we have defined this in Section 3). We can now make the following definition:

Definition 6.4. Let F be a holomorphic correspondence on $\mathbb{P}^1$, let $N \in \mathbb{Z}_+$, and let $w_0 \in \mathbb{P}^1$. Let $\boldsymbol{\mathcal{W}} := (w_0, w_{-1}, \dots, w_{-N}; \alpha_1, \dots, \alpha_N) \in \mathscr{P}_{-N}(F, w_0)$. We call the list $({}^\dagger F_{(\boldsymbol{\mathcal{W}},1)}, \dots, {}^\dagger F_{(\boldsymbol{\mathcal{W}},N)})$ a *regular branch of a backward iteration of F along $\boldsymbol{\mathcal{W}}$* if:

1) $w_0, w_{-1}, \dots, w_{-N+1}$ are regular values.
2) For each $j = 1, \dots, N$, ${}^\dagger F_{(\boldsymbol{\mathcal{W}},j)}$ is a holomorphic function defined by

$$ {}^\dagger F_{(\boldsymbol{\mathcal{W}},j)}(\zeta) \; := \; \pi_1 \circ \big(\, \pi_2|_{\mathcal{D}(\boldsymbol{\mathcal{W}},j)} \big)^{-1}(\zeta) \;\; \forall \zeta \in \pi_2(\mathcal{D}(\boldsymbol{\mathcal{W}}, j)),$$

where $\mathcal{D}(\boldsymbol{\mathcal{W}}, j)$ is a *local* irreducible component of $\Gamma^\bullet_{\alpha_j}$ at the point (w_{-j}, w_{-j+1}) such that: (a) $\mathcal{D}(\boldsymbol{\mathcal{W}}, j)$ is smooth; (b) π_2 restricted to $\mathcal{D}(\boldsymbol{\mathcal{W}}, j)$ is injective; and (c) $\pi_2(\mathcal{D}(\boldsymbol{\mathcal{W}}, j)) \supset \pi_1(\mathcal{D}(\boldsymbol{\mathcal{W}}, j-1))$ when $j \geq 2$.

The above definition is the analogue — for the scenario in which we are interested — of the notion of a "regular inverse branch of F of order N" introduced by Dinh in [8].

The following is the key proposition needed to prove Theorem 3.4.

Proposition 6.5. *Let F be a holomorphic correspondence of $\mathbb{P}^1$ having all the properties stated in Theorem 3.4 and let $\mathcal{R}$ be a strong repeller that is disjoint from the set of critical values of F. Then, there exists a closed set $B \subset \mathbb{P}^1$ such that $B^\circ \supset \mathcal{R}$ and such that:*

i) *The operator* $\mathbb{A}_B := d_1(F)^{-1}\,\mathbf{\Lambda}|_B$, *where* $\mathbf{\Lambda}|_B$ *is as defined in* (3.3) *with* B *replacing* Ω, *maps* $\mathcal{C}(B;\mathbb{C})$ *into itself.*

ii) *There exists a probability measure* $\mu_B \in \mathcal{C}(\mathbb{P}^1;\mathbb{R})^*$ *that satisfies* $\mu_B \circ \mathbb{A}_B = \mu_B$ *and such that*

$$\lim_{n\to\infty}\sup_B\left|\mathbb{A}^n[\varphi] - \int_B \varphi\,d\mu_B\right| \;=\; 0 \quad \forall \varphi \in \mathcal{C}(B;\mathbb{C}). \tag{6.1}$$

Proof. Let $a_0 \in \mathcal{R}$ and let U be an open neighbourhood of $\mathcal{R}$ such that:

- For each $w \in U$, there is a sequence $\{a_n(w)\}_{n\in\mathbb{Z}_+}$ such that $a_n(w) \in {}^{\dagger}F^n(w)$ for each n, and $a_n(w) \longrightarrow a_0$ as $n \to \infty$.
- U contains no critical values of F.

By Result 6.2 and Result 6.3, we can find an open neighbourhood W of $\mathcal{R}$ such that $\overline{W} \subset U$ and $\overline{W}$ is a closed attractor block for the relation $|{}^{\dagger}\Gamma|$.

Repeating the last argument once more, we can find a closed attractor block, B, for $|{}^{\dagger}\Gamma|$ such that

$$\mathcal{R} \subset B^{\circ} \subset B \subset W \Subset U.$$

By the above chain of inclusions and by the definition of the term "attractor block", it follows that the operator $\mathbb{A}_B$ maps $\mathcal{C}(B;\mathbb{C})$ into itself.

Claim 1. *For each fixed* $\varphi \in \mathcal{C}(B;\mathbb{C})$, $\{\mathbb{A}^n_B[\varphi]\}_{n\in\mathbb{Z}_+}$ *is an equicontinuous family.*

It is easy to see that $\mathcal{R}$ is a closed proper subset. We can thus make a useful observation:

> We can choose W so that $\mathbb{P}^1 \setminus \overline{W}$ is non-empty. Hence, we can choose coordinates in such a way that we may view W as lying in $\mathbb{C}$, and that $W \Subset \mathbb{C}$. We shall work with respect to these coordinates in the remainder of this proof.

Let us pick a point w_0 in B (which, by construction, is a regular value) and let $D(w_0)$ be a small disc centered at w_0 such that $\overline{D(w_0)} \subset W$. Let us fix an $N > 1$ and consider a path $\boldsymbol{\mathcal{W}} \in \mathscr{P}_{-N}(F, w_0)$. Recall that, by construction:

$${}^{\dagger}F(\overline{W}) \;\subset\; W. \tag{6.2}$$

We can infer from (6.2) that there exists a regular branch $({}^{\dagger}F_{(\boldsymbol{\mathcal{W}},1)}, \dots, {}^{\dagger}F_{(\boldsymbol{\mathcal{W}},N)})$ of a backward iteration of F along $\boldsymbol{\mathcal{W}}$. To see why, first note that, as w_0 is a regular value and $\overline{D(w_0)} \subset W$, we get:

(a_1) There is an open neighbourhood U_0 of w_0 containing only regular values.

(b_1) Writing

$$\widetilde{U}_1 := \text{the connected component of } \pi_2^{-1}(U_0) \cap \Gamma^{\bullet}_{\alpha_1} \text{ containing } (w_{-1}, w_0),$$

and defining $\mathcal{D}(\boldsymbol{\mathcal{W}}, 1) :=$ any one of the irreducible components of the (local) complex-analytic variety $\widetilde{U}_1$, the function

$${}^{\dagger}F_{(\boldsymbol{\mathcal{W}},1)} \;:=\; \pi_1 \circ \big(\,\pi_2|_{\mathcal{D}(\boldsymbol{\mathcal{W}},1)}\big)^{-1} \text{ is holomorphic on } U_0.$$

(c_1) The open set $U_1 := {}^{\dagger}F_{(\boldsymbol{\mathcal{W}},1)}(U_0) \subset W$ and hence contains only regular values.

The assertion (c_1) follows from the fact that $U_1 \subset {}^{\dagger}F(\overline{W}) \subset W$ and that the latter contains no critical values.

Let us now, for some $k \in \mathbb{Z}_+$, $k \leq N-1$, *assume* the truth of the statements (a_k), (b_k) and (c_k), which are obtained by replacing all the subscripts 0 and 1 in (a_1), (b_1) and (c_1) (except the subscript in π_1) by $k-1$ and k, respectively. Now, (a_{k+1}) follows from (c_k). Defining $\mathcal{D}(\boldsymbol{\mathcal{W}}, k+1)$ in exact analogy to $\mathcal{D}(\boldsymbol{\mathcal{W}}, 1)$, and writing

$$^{\dagger}F_{(\boldsymbol{\mathcal{W}},\,k+1)} \;:=\; \pi_1 \circ \big(\, \pi_2|_{\mathcal{D}(\boldsymbol{\mathcal{W}},\,k+1)}\,\big)^{-1},$$

the holomorphicity of $(\pi_2|_{\mathcal{D}(\boldsymbol{\mathcal{W}},\,k+1)})^{-1}$ follows from (a_{k+1}) and our definition of a regular value of F. Thus (b_{k+1}) holds true. We get (c_{k+1}) by appealing once again to (6.2) and using the fact that $U_k \subset W$. By induction, therefore, a regular branch of a backward iteration of F along $\boldsymbol{\mathcal{W}}$ exists.

We thus conclude the following:

- $^{\dagger}F_{(\boldsymbol{\mathcal{W}},\,N)} \circ \cdots \circ {}^{\dagger}F_{(\boldsymbol{\mathcal{W}},\,1)} \in \mathcal{O}(D(w_0))$; and
- $^{\dagger}F_{(\boldsymbol{\mathcal{W}},\,N)} \circ \cdots \circ {}^{\dagger}F_{(\boldsymbol{\mathcal{W}},\,1)}(D(w_0)) \subset W \Subset \mathbb{C}$.

Recall that $\boldsymbol{\mathcal{W}}$ was arbitrarily chosen from $\mathscr{P}_{-N}(F, w_0)$ and that the arguments in the last two paragraphs hold true for any choice of $\mathcal{D}(\boldsymbol{\mathcal{W}}, j)$, $1 \leq j \leq N$, and for any $N \in \mathbb{Z}_+$. Thus, by Montel's theorem, we infer the following important fact: the family

$$\begin{aligned} ^{\dagger}\mathscr{F}(w_0) \;:=\; \Big\{ & {}^{\dagger}F_{(\boldsymbol{\mathcal{W}},\,N)} \circ \cdots \circ {}^{\dagger}F_{(\boldsymbol{\mathcal{W}},\,1)} \in \mathcal{O}(D(w_0)) : \boldsymbol{\mathcal{W}} \in \mathscr{P}_{-N}(F, w_0) \textit{ for some } N \in \mathbb{Z}_+ \\ & \textit{\& } ({}^{\dagger}F_{(\boldsymbol{\mathcal{W}},\,N)}, \dots, {}^{\dagger}F_{(\boldsymbol{\mathcal{W}},\,1)}) \textit{ is a regular branch of a backward iteration of } F \Big\} \\ & \text{is a normal family.} \end{aligned} \tag{6.3}$$

Pick a $\varphi \in \mathcal{C}(B; \mathbb{C})$ and let $\varepsilon > 0$. As B is compact, there exists a number $\delta(\varepsilon) > 0$ such that:

$$|z_1 - z_2| < \delta(\varepsilon) \;\Rightarrow\; |\varphi(z_1) - \varphi(z_2)| < \varepsilon \quad \forall z_1, z_2 \in B. \tag{6.4}$$

We pick a $\zeta \in B$. By taking $\zeta = w_0$ in the discussion in the previous paragraph, we infer from the normality of the family $^{\dagger}\mathscr{F}(\zeta)$ that we can find a sufficiently small number $r(\varepsilon, \zeta) > 0$ such that:

$$|\xi - \zeta| < r(\varepsilon, \zeta) \text{ and } \xi \in B \;\Rightarrow\; |\psi(\xi) - \psi(\zeta)| < \delta(\varepsilon) \quad \forall \psi \in {}^{\dagger}\mathscr{F}(\zeta). \tag{6.5}$$

Now, for each $\zeta \in B$ write:

$$\mathscr{IB}(N, \zeta) \;:=\; \{(\psi, \boldsymbol{\mathcal{W}}) : \psi \text{ is some } {}^{\dagger}F_{(\boldsymbol{\mathcal{W}},\,N)} \circ \cdots \circ {}^{\dagger}F_{(\boldsymbol{\mathcal{W}},\,1)} \text{ in } {}^{\dagger}\mathscr{F}(\zeta),\; \boldsymbol{\mathcal{W}} \in \mathscr{P}_{-N}(F, \zeta)\}.$$

If $(\zeta_0, \zeta_{-1}, \dots, \zeta_{-N}; \boldsymbol{\alpha}) =: \boldsymbol{\mathcal{W}}$ is a path of backward iteration (with ζ_0 being the above ζ), basic intersection theory tells us that the local intersection multiplicity of $\Gamma^{\bullet}_{\alpha_j}$ with $\mathbb{P}^1 \times \{\zeta_{-j+1}\}$ at $(\zeta_{-j}, \zeta_{-j+1})$ equals the number of distinct branches $^{\dagger}F_{(\boldsymbol{\mathcal{W}},j)}$ one can construct according to the above inductive prescription (this number is greater than 1 if $\Gamma^{\bullet}_{\alpha_j}$ has a normal-crossing singularity at $(\zeta_{-j}, \zeta_{-j+1})$). From this, and from the iterative construction of the $^{\dagger}F_{(\boldsymbol{\mathcal{W}},\,N)}$'s above, it follows that:

$$\mathsf{Card}[\,\mathscr{IB}(N, \zeta)\,] \;=\; d_1(F)^N \quad \forall \zeta \in B. \tag{6.6}$$

From (6.4), (6.5) and (6.6), we get:

$$\begin{aligned}|\mathbb{A}^n_B[\varphi](\xi) - \mathbb{A}^n_B[\varphi](\zeta)| \;&=\; \Bigg|\sum_{(\psi,\boldsymbol{\mathcal{W}})\in\mathscr{IB}(n,\zeta)} d_1(F)^{-n}(\varphi\circ\psi(\zeta) - \varphi\circ\psi(\xi))\Bigg| \\ &\leq d_1(F)^{-n}\sum_{(\psi,\boldsymbol{\mathcal{W}})\in\mathscr{IB}(n,\zeta)} |\varphi\circ\psi(\zeta) - \varphi\circ\psi(\xi)| \;<\; \varepsilon \\ &\qquad \forall \xi\in B \text{ such that } |\xi-\zeta| < r(\varepsilon,\xi) \text{ and } \forall n\in\mathbb{Z}_+.\end{aligned}$$

The above holds true for each $\zeta\in B$. This establishes Claim 1.

In what follows, the term *unitary spectrum* of an operator on a complex Banach space will mean the set of all eigenvalues of the operator of modulus 1, which we will denote by $\mathsf{Spec}_{\mathsf{U}}$.

Claim 2: $\mathsf{Spec}_{\mathsf{U}}(\mathbb{A}_B) = \{1\}$, *and the eigenspace associated with* 1 *is* $\mathbb{C}$.
The proof of the above claim is essentially the one in [14]. However, to make clear the role that the properties of $\mathcal{R}$ play, we repeat most of the details of Lyubich's argument. Let us fix a $\lambda\in\mathsf{Spec}_{\mathsf{U}}(\mathbb{A}_B)$ and let $\varphi_\lambda\in\mathcal{C}(B;\mathbb{C})$ be an associated eigenfunction. Let $\zeta_*\in B$ be such that $|\varphi_\lambda(\zeta_*)| = \max_B|\varphi_\lambda|$. By definition

$$d_1(F)^{-1}\sum_{(\psi,\boldsymbol{\mathcal{W}})\in\mathscr{IB}(1,\zeta_*)} \varphi_\lambda\circ\psi(\zeta_*) \;=\; \lambda\varphi_\lambda(\zeta_*). \tag{6.7}$$

The above equality would fail if, for some ψ occurring above, $|\varphi_\lambda\circ\psi(\zeta_*)| < |\varphi_\lambda(\zeta_*)|$. Furthermore, if, for some ψ occurring above, $\varphi_\lambda\circ\psi(\zeta_*)$ and $\lambda\varphi_\lambda(\zeta_*)$ do not lie on the same ray through $0\in\mathbb{C}$, then cancellations would lead to (recall the fact (6.6) and that $|\varphi_\lambda\circ\psi(\zeta_*)| = |\varphi_\lambda(\zeta_*)|$)

$$\Bigg|\sum_{(\psi,\boldsymbol{\mathcal{W}})\in\mathscr{IB}(1,\zeta_*)} \varphi_\lambda\circ\psi(\zeta_*)\Bigg| \;<\; d_1(F)|\varphi_\lambda(\zeta_*)|,$$

which is a contradiction. Thus $\varphi_\lambda(x) = \lambda\varphi_\lambda(\zeta_*)\ \forall x\in {}^{\dagger}F(\zeta_*)$. Iterating, we get

$$\varphi_\lambda(x) \;=\; \lambda^n\varphi_\lambda(\zeta_*) \quad \forall x\in {}^{\dagger}F^n(\zeta_*),\ \ n=1,2,3,\dots \tag{6.8}$$

Since, by construction, $B\subset U$, there exists an $x_n\in {}^{\dagger}F^n(\zeta_*)$, $n=1,2,3,\dots$, such that $x_n\longrightarrow a_0$. Therefore, owing to (6.8), the sequence $\{\lambda^n\varphi_\lambda(\zeta_*)\}_{n\in\mathbb{Z}_+}$ is a convergent sequence. As $\varphi_\lambda\not\equiv 0$ (by definition), this implies that $\lambda = 1$.

Observe that, having determined that $\lambda = 1$, (6.8) also gives

$$\varphi_\lambda(\zeta_*) \;=\; \varphi_\lambda(a_0). \tag{6.9}$$

Note that $\overline{\mathbb{A}[\varphi]} = \mathbb{A}[\overline{\varphi}]\ \forall\varphi\in\mathcal{C}(B;\mathbb{C})$. Hence, $\mathsf{Re}\varphi_\lambda$ and $\mathsf{Im}\varphi_\lambda$ are also eigenvectors of $\mathbb{A}$ associated to $\lambda = 1$. Thus, we have the following analogue of (6.7):

$$d_1(F)^{-1}\sum_{(\psi,\boldsymbol{\mathcal{W}})\in\mathscr{IB}(1,z^\bullet)} \mathsf{Re}\varphi_\lambda\circ\psi(z^\bullet) \;=\; \lambda\mathsf{Re}\varphi_\lambda(z^\bullet),$$

where $z^\bullet\in B$ stands for either a point of global maximum or a point of global minimum of $\mathsf{Re}\varphi_\lambda$. Using the above as a starting point instead of (6.7) and repeating, with appropriate modifications, the argument that begins with (6.7) and ends at (6.9), we get:

$$\min_B(\mathsf{Re}\varphi_\lambda) \;=\; \mathsf{Re}\varphi_\lambda(a_0) \;=\; \max_B(\mathsf{Re}\varphi_\lambda). \tag{6.10}$$

Similarly, we deduce that:

$$\min_B(\mathsf{Im}\varphi_\lambda) \;=\; \mathsf{Im}\varphi_\lambda(a_0) \;=\; \max_B(\mathsf{Im}\varphi_\lambda).$$

Combining the above with (6.10), we conclude that, for any eigenvector φ_λ associated to $\lambda = 1$, $\varphi_\lambda \equiv$ *constant*. This establishes Claim 2.

To complete this proof, we need the following:

Result 6.6 (Lyubich, [14])**.** *Let $\mathscr{B}$ be a complex Banach space. Let $A : \mathscr{B} \longrightarrow \mathscr{B}$ be a linear operator such that $\{A^n(v)\}_{n\in\mathbb{Z}_+}$ is a relatively-compact subset of $\mathscr{B}$ for each $v \in \mathscr{B}$. Assume that $\mathsf{Spec}_{\mathsf{U}}(A) = \{1\}$ and that 1 is a simple eigenvalue. Let $h \neq 0$ be an invariant vector of A. Then, there exists a linear functional μ that satisfies $\mu \circ A = \mu$ and $\mu(h) = 1$, and such that*

$$A^n(v) \longrightarrow \mu(v)h \;\; \text{as } n \to \infty$$

for each $v \in \mathscr{B}$.

Take $\mathscr{B} = \mathcal{C}(B;\mathbb{C})$ in the above theorem. Note that $|\mathbb{A}^n_B[\varphi]|$ is bounded by $\max_B|\varphi|$ for $n = 1, 2, 3, \dots$ Thus, in view of Claims 1 and 2, $\mathbb{A}_B$ satisfies all the hypotheses of Result 6.6. Hence (recall that the function that is identically 1 on B is an eigenvector of $\mathbb{A}_B$) there is a regular complex Borel measure μ_B on B such that $\int_B 1\, d\mu_B = 1$, and

$$\lim_{n\to\infty}\sup_B\left|\mathbb{A}^n[\varphi] - \int_B \varphi\, d\mu_B\right| \;=\; 0 \quad \forall\varphi \in \mathcal{C}(B;\mathbb{C}).$$

It is clear from the above equation that μ_B is a positive measure. Hence it is a probability measure on B. ☐

The proof of Theorem 3.4. Let B be any closed set having the properties listed in the conclusion of Proposition 6.5. Let μ_B be the probability measure associated to this B. We claim that μ_F is given by defining:

$$\int_{\mathbb{P}^1}\varphi\, d\mu_F \;:=\; \int_B \varphi|_B\, d\mu_B \quad \forall\varphi \in \mathcal{C}(\mathbb{P}^1;\mathbb{C}).$$

We must show that μ_F does not depend on the choice of B. The proof of this is exactly as given in [14, Theorem 1]. We fix a point $z \in \mathcal{R}$. So, $z \in B$ for any choice of B. Thus,

$$\begin{aligned}\int_{\mathbb{P}^1}\varphi\, d\mu_F \;&=\; \lim_{n\to\infty}\mathbb{A}^n_B[\varphi](z) && \text{(by Proposition 6.5)}\\ &=\; \lim_{n\to\infty} d_1(F)^{-n}\mathbf{\Lambda}^n[\varphi](z), && (\text{since } {}^{\dagger}F^n(z) \subset {}^{\dagger}F^n(\mathcal{R}) \subset \mathcal{R})\end{aligned}$$

where $\mathbf{\Lambda}$ is as described in the passage following (3.3). The last line is independent of B. Hence the claim.

By the above calculation, we also see that $\int_{\mathbb{P}^1} 1\, d\mu_F = 1$. Thus, μ_F is a probability measure.

Let U be the open set described at the beginning of the proof of Proposition 6.5. We now define:

$$\mathfrak{B}(\mathcal{R}) := \Big\{B \subsetneq U : B \text{ is closed, } B^\circ \supset \mathcal{R},\ {}^{\dagger}F(B) \subset B \text{ and there exists a closed attractor block } B_* \text{ for } |{}^{\dagger}\Gamma| \text{ s.t. } B \subset B_*^\circ \subset B_* \subset U\Big\}.$$

We see from the proof of Proposition 6.5 that, owing to our hypotheses, $\mathfrak{B}$ is non-empty. Hence

$$U(F,\mathcal{R}) := \bigcup\nolimits_{B\in\mathfrak{B}(\mathcal{R})} B^\circ$$

is a non-empty open set that contains $\mathcal{R}$. Let $z \in U(F,\mathcal{R})$. There exists a $B \in \mathfrak{B}$ such that $z \in B^\circ$. A close look at the essential features of its proof reveals that this B has all the properties listed in the conclusion of Proposition 6.5. Consider any $\varphi \in \mathcal{C}(\mathbb{P}^1;\mathbb{C})$. We now apply Lemma 4.3 to get

$$\begin{aligned} d_1(F)^{-n}\langle (F^n)^*(\delta_z),\varphi\rangle &= d_1(F)^{-n}\mathbf{\Lambda}^n[\varphi](z) && \text{(from (3.3) and Lemma 4.3)}\\ &= \mathbb{A}_B^n[\varphi](z) && (\text{since } {}^{\dagger}F^n(z) \subset {}^{\dagger}F^n(B^\circ) \subset B)\\ &\longrightarrow \int_{\mathbb{P}^1} \varphi\, d\mu_F \ \text{ as } n\to\infty, && (6.11) \end{aligned}$$

and this holds true *for any* $z \in U(F,\mathcal{R})$. The last line follows from our observations above on μ_F. Now note that, by construction, ${}^{\dagger}F(z) \subset U(F,\mathcal{R})$ for each $z \in U(F,\mathcal{R})$. Therefore, in view of equation (3.3), it follows from (6.11) that $F^*(\mu_F) = d_1(F)\mu_F$.

The proof of the fact that μ_F is invariant under F is the same as in the proof of Theorem 3.1. □

7. An example

The purpose of this section is to show that it is easy to construct examples of holomorphic correspondences on $\mathbb{P}^1$ that satisfy all the conditions stated in Theorem 3.4, but, unlike the example discussed in Remark 3.5, have "large" repellers.

Choose a complex polynomial p with $\deg(p) \geq 2$ such that its Julia set $\mathcal{J}_p \subsetneq \mathbb{P}^1$ and such that no critical values of p lie in $\mathcal{J}_p$. It follows — see, for instance, [14, §4] — that there is a compact set B such that $B^\circ \supset \mathcal{J}_p$ and avoids the critical values of p, and such that $p^{-1}(B) \subset B^\circ$. Next, choose a polynomial Q with $\deg(Q) \geq 2$ and having the following properties:

a) Q has an attractive fixed point, call it ζ_0, in $\mathcal{J}_p$;
b) B lies in the basin of attraction (under the action of Q) of ζ_0.

In view of (*b*), we can find a positive integer N that is so large that

- $Q^N(B) \subset B^\circ$;
- $N\deg(Q) \geq \deg(p)$.

Let us write $q := Q^N$.

Next, we define:

$$\begin{aligned}\Gamma_1 &:= \{([z_0:z_1],[w_0:w_1]) : w_0^{\deg(q)} z_1 - w_0^{\deg(q)} z_0\, q(w_1/w_0) = 0\},\\ \Gamma_2 &:= \{([z_0:z_1],[w_0:w_1]) : w_1 z_0^{\deg(p)} - w_0 z_0^{\deg(p)} p(z_1/z_0) = 0\}.\end{aligned}$$

The projective coordinates are so taken that $[0:1]$ stands for the point at infinity.

We set $\mathit{\Gamma} := \Gamma_1 + \Gamma_2$ and let F denote the correspondence determined by $\mathit{\Gamma}$. Clearly $d_0(F) = \deg(q) + 1 \geq \deg(p) + 1 = d_1(F)$. By construction, ${}^{\dagger}F(B) \subset B^{\circ}$. In other words, B an attractor block for the relation $|{}^{\dagger}\mathit{\Gamma}|$. Therefore, it follows from Result 6.2 and Definition 3.2 that F has a repeller $\mathcal{R} \subset B^{\circ}$; this repeller is just $\omega(B; |{}^{\dagger}\mathit{\Gamma}|)$.

It is easy to see that

$$\bigcap_{n\geq 0} \overline{\bigcup_{k\geq n} {}^{\dagger}F^k(B)} \subseteq \omega(B; |{}^{\dagger}\mathit{\Gamma}|)$$

(the reader may look up [15, Theorem 5.1] for a proof). The above implies that

$$\mathcal{J}_p \subseteq \mathcal{R} \subseteq B^{\circ}.$$

Thus, the correspondence F defined above has a repeller that is disjoint from the set of critical values of F. Owing to (a) and (b) above, for each $w \in B^{\circ}$, there exists a point $a_n(w) \in {}^{\dagger}F^n(w)$ such that $a_n(w) \longrightarrow \zeta_0$. Hence, $\mathcal{R}$ is a strong repeller.

Acknowledgement. A part of this work was carried out during a visit by Shrihari Sridharan to the Indian Institute of Science. He would like to thank the Department of Mathematics, Indian Institute of Science, for its support and hospitality.

REFERENCES

[1] H. Brolin, *Invariant sets under iteration of rational functions*, Ark. Mat. **6** (1965) 103-14.

[2] S. Bullett, *Dynamics of quadratic correspondences*, Nonlinearity **1** (1988), no. 1, 27-45.

[3] S. Bullett, *A combination theorem for covering correspondences and an application to mating polynomial maps with Kleinian groups*, Conform. Geom. Dyn. **4** (2000), 75-96.

[4] S. Bullett and M. Freiberger, *Holomorphic correspondences mating Chebyshev-like maps with Hecke groups*, Ergodic Theory Dynam. Systems **25** (2005), no. 4, 1057-1090.

[5] S. Bullett, C. Penrose, *Mating quadratic maps with the modular group*, Invent. Math. **115** (1994), no. 3, 483-511.

[6] S. Bullett and C. Penrose, *Regular and limit sets for holomorphic correspondences*, Fund. Math. **167** (2001), no. 2, 111-171.

[7] J.B. Conway, *Functions of One Complex Variable*, Springer-Verlag, New York, 1973.

[8] T.-C. Dinh, *Distribution des préimages et des points périodiques d'une correspondance polynomiale*, Bull. Soc. Math. France **133** (2005), no. 3, 363-394.

[9] T.-C. Dinh and N. Sibony, *Distribution des valeurs de transformations méromorphes et applications*, Comment. Math. Helv. **81** (2006), no. 1, 221-25.

[10] T.-C. Dinh and N. Sibony, *Upper bound for the topological entropy of a meromorphic correspondence*, Israel J. Math. **163** (2008), 29-44.

[11] A. Douady, *Systèmes dynamiques holomorphes*, Séminaire Bourbaki, Vol. 1982/83, 39-63, Astérisque **105-106**, 1983.

[12] F. Kirwan, *Complex Algebraic Curves*, London Mathematical Society Student Texts 23, Cambridge University Press, Cambridge, 1992.

[13] T. Lei, *Matings of quadratic polynomials*, Ergodic Theory Dynam. Systems **12** (1992), no. 3, 589-620.

[14] M. Ju. Ljubich, *Entropy properties of rational endomorphisms of the Riemann sphere*, Ergodic Theory Dynam. Systems **3** (1983), no. 3, 351-385.

[15] R. McGehee, *Attractors for closed relations on compact Hausdorff spaces*, Indiana Univ. Math. J. **41** (1992), no. 4, 1165-1209.

[16] A. Russakovskii and B. Shiffman, *Value distribution for sequences of rational mappings and complex dynamics*, Indiana Univ. Math. J. **46** (1997), no. 3, 897-932.

[17] I.R. Shafarevich, *Basic Algebraic Geometry 1, Varieties in Projective Space* (2nd ed.; translated from the Russian & with notes by Miles Reid), Springer-Verlag, Berlin, 1994.

Department of Mathematics, Indian Institute of Science, Bangalore 560012, India
E-mail address: `bharali@math.iisc.ernet.in`

Chennai Mathematical Institute, Chennai 603103, India
E-mail address: `shrihari@cmi.ac.in`